

Index

Page numbers in *italic* denote figures. Page numbers in **bold** denote tables.

- Acastidae, 305
Acastidae Delo, 304–305
acritarchs, 365
 cluster analysis, 387
Acrocephalitidae, 325
actinocerids, 433, 437, 440
adaptive radiation of vertebrates in Ordovician, 451–452
 biogeography of Ordovician vertebrates, 451–452
 climatic context of Ordovician vertebrates, 452
 palaeobiogeography of Early Palaeozoic vertebrates, 451–452
Aeronian cluster and ordination tests, 209
Aeronian-Telychian, 209–210
Afghanistan palaeogeographical units, 279
Afghanodesmatids Ordovician distributions, 224
agnathans, 449, 455
Alai terrane
 fossils, 279
 trilobite, 283–284
Alaskan areas, 213–214
Alsatspididae, 320
Alsatspididae Turner, 319–320
Altai-Sayan area, palaeogeographical units, 276
Ambonychids, Ordovician distributions, 225
Anatolepis, 451
Annamia, palaeogeographical reconstructions, 8–9
Anomalodesmatans, 223
Antarctica, palaeogeographical units, 281
Anti-Atlas Mountains trilobites, 285
Antipleuridae, 231
Archaeocyatha, 59–64
 bioconstruction distribution, 61
 biostratigraphy, 59
 Cambrian biostratigraphy and biogeography, 59–64
 cluster analysis, 62
 database and website, 63–64
 global palaeogeographical distribution, 61
 palaeoecology, 60–62
 palaeogeography, 62–63
 radiocyathan genera, 62
 systematic studies history, 59
archaeocyaths v. trilobites, **60**
archipelagos, 26
Arctic Alaska-Chukotka Microcontinent, 7
 palaeogeographical reconstructions, 9
Arenig Stage bivalves, 223
Argentina
 eocrinoids, 180
 Famatina System, 277
Argentinean Precordillera, 216
Arhourilla, 222
Armorian Terrane Assemblage, 12
 palaeogeographical reconstructions, 9–10
Arthropod genera, 38
Asaphidae, 317
Asaphida Salter, 317–321
 Family Alsatspididae Turner, 319–320
 Family Ceratopygidae Linnarsson, 318
 Family Dionididae Gurich, 320
 Family Nileidae Angelin, 319
 Family Raphiophoridae Angelin, 320
 Family Trinucleidae Hawle & Corda, 320–321
 Superfamily Asaphoidea Burmeister, 317–318
 Superfamily Cyclopygoidea Raymond, 318–319
 Superfamily Trinucleoidea Hawle & Corda, 319–321
 synopsis of Ordovician trilobite distribution and diversity,
 317–321
Asaphoidea Burmeister, 317–318
Ashgill units within Continental Margin Area
 coral genera, **102**
Asteroids, 178–179
Asterozoans, 177–179
 Asteroids, 178–179
 Ophiuroids, 178
 palaeobiogeography of Ordovician echinoderms, 177–179
 Somasteroids, 177–178
Astropolichnus hispanicus (Crimes, Legg, Marcos and Arboleya), 45
Atdabanian Stage, palaeogeographical distribution, 61
Aulacopleurida, 311
Aulacopleurida Adrain, 312–317
 Family Aulacopleuridae Angelin, 312–313
 Family Bathyuridae Walcott, 313–314
 Family Brachymetopidae Prantl & Přibyl, 314
 Family Dimeropygidae Hupe, 314
 Family Holotrachelidae Warburg, 314–315
 Family Hystricuridae Hupe, 315
 Family Rorringtoniidae Owens in Owens & Hamann, 315–316
 Family Scharysiidae Osmólska, 316
 Family Telephinidae Marek, 316–317
 synopsis of Ordovician trilobite distribution and diversity, 312–317
Aulacopleuridae
 global taxonomic richness, 313
Aulacopleuridae Angelin, 312–313
Australia
 bryozoan faunas, 147
 genera, 203
 GMM [Gastropods, monoplacophorans and mimospirids], 215
 mid-Ordovician faunas, 226
 palaeogeographical units, 281
 supercontinent breakup, 281
Avalonia, 277
 blastozoans, 181
 cephalopods, 433
 palaeogeographical reconstructions, 10
 Silurian, 232–234
 Trilobites, Cambrian Epoch 3, 286
 West Gondwana, 282
axonophorans, 424
Baltica, 268, 351
 biogeographical analysis of trilobites, 287
 bryozoan, 146
 cephalopods, 433–435
 echinoderm assemblage, 191
 GMM [Gastropods, monoplacophorans and mimospirids], 214–215
 Heterococonchs *Lyrodesma poststriatum*, 229
 Ordovician, 369
 Ordovician echinoderm assemblages, 174
 Ordovician echinoderms, 186
 palaeogeographical province, 186
 palaeogeographical reconstructions, 8
 palaeogeographical units, 274
 Silurian, 232–234
Baltica-Siberia Subrealm, 452
Baltoscandia, 337
 Ordovician facies zones, 338
basslerocerids, 435
Bathmoceras, 432
Bathycheilidae, 309
Bathycheilidae Přibyl, 309
bathypelagic species, 362
Bathyuridae, 313
Bathyuridae Walcott, 313–314
Batostoma, 148
Bavarillidae, 310
Bavarillidae Sdzuy, 309
bellerophonoid gastropods, 200
bellerophonoid *Peelerophon*, 200
Benbolt Formation, 180

- benthic faunas, 16
benthic species
 stratigraphical distribution, 357
 testing Silurian palaeogeography, 362–363
benthic trilobites, 46
Beothuka terranova, 411
Beyrichia admixta, 358
bilaterians, 40
binodicipes, 356
biodiversity Ordovician, 127–142
biofacies, 376
biogeographers, 26–27
biogeographical patterns
 Early Palaeozoic Rostroconchia, 243–260
 Ordovician ostracods, 337–351
biogeographical provinces, 25
biogeographical regions, **26**
 ranks, **31**
biogeography
 bryozoans, Early Palaeozoic, 145–154
 Early Late Ordovician, 350
 early to mid Palaeozoic marine phytoplankton, 365–389
 Ordovician, 127–142
 Ordovician and Silurian, 199–217
 Ordovician and Silurian GMM [Gastropods, monoplacophorans and mimospirids], 199–217
 Ordovician and Silurian Stromatoporoidea, 67–78
 Ordovician distribution, 67–73
 Ordovician linguliform and craniiform brachiopods, 117–124
 Silurian distribution, 73–77
Bioherms, 109
biosedimentology reefs, 62
biostratigraphy of Archaeocyatha, 59
biostromes, 107
biozones of trilobites and archaeocyaths, **60**
biseriate diversification, 423
bivalves, 221–236
 Arenig Stage, 223
 Cambrian, 221–222
 Lower Palaeozoic palaeobiogeography, 221–236
 Ordovician, 222–230
 Ordovician diversification, 222
 Silurian, 230–235
blastozoans
 Avalonian Province, 181
 Furongian, 162
 Ordovician echinoderms palaeobiogeography, 179–184
 Sandbian-Katian interval, 181
Boda event, 72
Bohemia palaeogeographical units, 278
Bolivia eocrinoids, 180
Botomian Stage
 Cambrian genera, 64
 global palaeogeographical distribution, 61
brachiopod diversification, 130
 phylogeography during Ordovician, 136–142
brachiopods, 117–124
 biogeography of Ordovician, 117–124
 craniiform, 121–122
 Laurentia, 134
 linguliform, 117–121
Brachymetopidae, 314
Brachymetopidae Prantl & Přibyl, 314
Brevilamnulella, 135
Britain Heteroconchs *Lyrodosma poststriatum*, 229–230
Bruno–Silesia palaeogeographical reconstructions, 12–13
bryozoan, 145–154
 Baltic Province, 146
 Dapingian, 146
 Dariwillian, 147
 Darrivilian, 146
 Early Palaeozoic biogeography, 145–154
 Floian, 146
 Hirnantian, 149, 149
 Katian, 146–149, 148
 Llandovery, 150–151, 150
 Ludlovian, 152
 Ludlow, 152–154
 Ordovician, 145–146
 Přídolí, 153, 154
 Sandbian, 146, 147
 Silurian, 149–150
 Tremadocian, 146
 Wenlock, 151–152, 151
Builth Inlier, 85
Burgess Shale, 40
 deposits, 35–40
 faunas, 81
Calcareans, 91
Calymenidae, 310
Calymenidae Swinnerton, 309–310
Calymenina Swinnerton, 309–310
Cambrian
 and archaeocyathan distribution, 61
 Archaeocyatha role, 59–64
 biostratigraphy and biogeography, 59–64
 Bivalvia, 221–222
 Botomian, 64
 Burgess Shale-type faunas, 81
 data cluster analysis, 371
 echinoderms, 160, 161
 global chronostratigraphical series and stages, 159
 lichakephalids, 301
 Major Early Palaeozoic events, 14–16
 non-stromatoporoid Porifera, 81–93, 86–91
 palaeobiogeographical units, 282
 palaeogeographical positions, **37**
 palaeogeographical reconstruction, 275
 Paleodictyon occurrences, 49–50
 phytoplankton, 366–367
 radiation, 39
 sponge palaeobiogeography, 83–84
 sponge distributions, 84
 tectonostratigraphic units, 280
 terrane or microcontinents, biogeographical concepts, 282
Cambrian clades
 analyses, 36
 clade dynamics at genus level, 37–39
 clade dynamics at species level, 36–37
 database, 35
 geographical ranges of genera, 37–38
 geographical ranges of species, 37
 groups studied, 36
 regions occupied v. temporal boundaries, 39
 spatio-temporal bias, 36
 temporal persistence of genera, 38–39
 temporal persistence of species, 37
Cambrian echinoderms, 157–167
 Cincta, 162–163
 Ctenocystoidea, 163
 diversity, 159–164, 166
 Edrioasteroidea, 159–161
 Eocrinoids, 161–162
 Helicoplacoidea, 159
 palaeobiogeographical patterns, 165–167
 palaeobiogeography, 157–167, 159–164
 palaeogeographical distribution, 159–164
 palaeogeographical distribution of Cambrian echinoderms, 159–164
 palaeogeographical units and chronostratigraphical nomenclature, 158–159
 record of disarticulated echinoderm ossicles, 164–165
 Rhombifera, 162
 Soluta, 163
 Stylophora, 163–164
Cambrian Epoch and geo-dispersal analysis, 285
Cambrian Epoch 2
 Furongian trilobite genera, 288
 trilobite, 285–286
 trilobite biogeography, 284

- Cambrian Epoch 3
 Avalonian trilobites, 286
 Furongian trilobite genera, 288
 PAE [Parsimony Analysis of Endemicity] analysis, 289
- Cambrian Explosion, 1
- Cambrian faunas, 166
- Cambrian Ordovician cephalopod palaeogeography and diversity, 429–444
 changing endemicity and beta-diversity, 438–439
 changing palaeogeographical pattern, 439–440
 diversity trends interpretation, 441–443
 general pattern, 438–441
 Late Ordovician, 443
 Mid Ordovician, 442–443
 palaeogeographical regions, 432–438
 regional diversity trends, 440
 widespread genera, 440–441
- Cambrian Series 2
 formations and species, 167
- Cambrian Series 3, 160
 Eocrinoid faunas, 162
- Cambrian Series 5
 formations and species, 167
- Cambrian Stage 5
 radiate echinoderms, 165
- Cambrian strata
 Moroccan Atlas, 278
 Tarim, 281
- Cambrian trace fossils
Astropolichnus hispanicus (Crimes, Legg, Marcos and Arboleya), 45
 Cambrian *Paleodictyon* occurrences, 49–50
Climactichnites Logan, 45–46
Cruziana d'Orbigny, 46
Dactyloidites Hall, 46–47
Didymaulichnus miettensis Young, 47
Oldhamia Forbes, 47–49
Paleodictyon meneghini, 49–50
 pre-trilobitic *Rusophycus* Hall, 50–52
Syringomorpha nilsoni (Torell), 52–53
- Cambrian trilobite, 273–290
 biogeographical links during Cambrian Epoch 2, 285–286
 biogeographical links during Cambrian Epoch 3, 286–287
 biogeography and timing of Cambrian radiation, 284–285
 Furongian biogeographical links, 287–289
 palaeobiogeography assessed using PAE [Parsimony Analysis of Endemicity], 273–290
- Campanario Formation, 52
- Canya, 222
- carbonates, 109
 mudmound facies pre-Hirnantian Ashgill, 135
- Carboniferous terranes, 11
- Cardiolidae, 231
- Catamarcaia*, 223
- Catillicephalidae, 325
- Catillicephalidae Raymond, 326
- Cellon Profile, 345
- Celtic group of brachiopod faunas, 132
- Celtic province, 132
- Central American terranes
 palaeogeographical reconstructions, 11
 palaeogeographical units, 277
- Central Asia
 cephalopods, 435
 palaeogeographical units, 279
- cephalic fields, 453
- cephalochordate, 450
- cephalopods
 Avalonia, Perunica, 433
 Baltica, 433–435
 Central Asia and organic belts, 435
 Eastern Gondwana and Sibumasu, 435
 global endemicity, **438**
 Laurentia, 435–436
 limestone biofacies, 231
 North China platform, 436–437
 Ordovician palaeogeographical regions, 432–438
 palaeogeography and diversity, 429–444
 Siberia and Far Russian East, 437
 South China platform, 437–438
 Tarim, 438
 Western Gondwana and Precordillera, 438
- Ceratopygidae, 318
- Ceratopygidae Linnarsson, 318
- Chazyan, 68
- Cheiruridae, 308
- Cheiruridae Hawle & Corda, 306
- Cheirurina Harrington & Leanza, 306–309
- Chinese tectonostratigraphic units, 283
- chitinozoans zooplankton maps, 400–401
- Chlorophyta, 37
- Cincinnati, 99
 biogeographical divisions of Laurentia, **99**
 Continental Margin Area, **102**
 coral genera, 97–112, **99**, **102**
- Cinctans
 Cambrian echinoderms, 162–163
 evolution, 163
- clades. *See also* Cambrian clades
 genus level diversity, **39**
 genus level spatio-temporal occurrence data, **39**
 species level diversity, **38**
 species level spatio-temporal occurrence data, **38**
- Clarkeia* Fauna, 7
- Clarkoceras*
 occurrences, 440
- Climactichnites* Logan, 45–46
- climograph, 27
- Clitambonitoidea, 138
- cluster analysis, 120
 acritarch assemblage, 387
 Aeronian-Telychian, 209–210
 archaeocyathan and radiocyathan genera, 62
 biogeography of Ordovician and Silurian GMM [Gastropods, monoplacophoran and mimospirids], 203–213
 Cambrian data, 371
 Dapingian-Darriwilian, 204–205
 Darriwilian-Sandbian, 205–207
 Floian time slice, 132
 Floian-Dapingian, 204
 Gorstian-Ludfordian, 212–213
 Hirnantian-Rhuddanian, 208–209
 Hirnantian time slice, 140
 Homerian-Gorstian, 211–212
 Katian-Hirnantian, 208
 Ludfordian-Přídolí, 213
 and ordination tests, 203–213
 Ordovician ostracod faunas, 350
 Rhuddanian-Aeronian, 209
 Sandbian-Katian, 207–208
 Sandbian time slice, 136
 Sheinwoodian-Homerian, 211
 Telychian-Sheinwoodian, 210
 Tremadocian-Floian, 203–204
 Tremadocian time slice, 130
- Coelenterata, 59
- Conocardiida, 244
- Conocardiida Neumayr, 243
- conocardiid rostroconchs geographical distribution, 244
- conodonts, 449
- continental blocks and terranes, 281
- continental crust, **28**
- continental disparity, 130
- continental drift theory, 28
- Continental Margin Area, 103, 109–111
 characteristic corals, 110
 coral biogeography in Late Ordovician (Cincinnati) of Laurentia, 109–111
 corals, 110
 duration and location, 109
 palaeogeography in tectonics, 29
- Coolibah Formation, 385

- coral genera
 biogeographical divisions in Laurentia, 98
 biogeography in Late Ordovician of Laurentia, 97–112
 Cincinnati, 97–112, **99**
 Continental Margin Area, **102**, 109–111, *110*
 Edgewood Province, **101**, 107–109, *108*
 Late Ordovician, 98
 Red River-Stone Mountain Province, 99–104, **100**, *103*
 Richmond Province, **101**, 105–107, *105*
- Cordilleran terranes, 8
 palaeogeographical reconstructions, 10
- Corynexochida Kobayashi, 299–301
 Family Illaenidae Hawle & Corda, 299
 Family Leiostegiidae Bradley, 300–301
 Family Missisquoiidae Hupe, 301
 Family Panderiidae Bruton, 299
 Family Styginidae Vogdes, 299–300
 synopsis of Ordovician trilobite distribution and diversity, 299–301
- Coryphidium bohemicum*, 367
- Cosmopolitan acritarchs, 383
- cosmopolitanism, **31**
- cothurnocystid cornutes, 164
- craniides, 121
 geographical distribution of Early Ordovician, *120*
- Craniiform brachiopods, 121–122
 biogeography, 117–124
 Early to Mid-Ordovician, 121
 Late Ordovician, 121–122
 Ordovician linguliform, 121–122
- craton (cratonic shield), **28**
- Cretaceous-style greenhouse, 403
- crinoids
 morphology, *184*
 Ordovician echinoderms palaeobiogeography, 184–185
 Tremadoc Series, 184
- Croizat, Leon, 26
- crustal fragments, 6
- Cruziana* d'Orbigny, 46
- cryptospores, 465
- cryptostome bryozoan, 145
- Ctenocystoidea, Cambrian echinoderms, 163
- Ctenocystoids Laurentia, 163
- Cuyania* trilobite, 283
- Cybelinae, 307
- Cyclocystoidea Echinozoans, 187
- Cyclopygidae, *318*
- Cyclopygoidea Raymond, 318–319
- Cyrtodontids Ordovician distributions, 224–225
- Czech Republic eocrinoids, 180
- Dactyloidites*
 distribution, *47*
miettensis distribution, *49*
- Dalmanitidae, *305*
- Dalmanitidae Vogdes, 305
- Dapingian Age, 418
 bryozoans, 146
 cluster analysis, *134*
 cluster and ordination tests, 204–205
 time slices, 130–133, *133*
- Dapingian Stage edrioasteroids, 187
- Darriwilian
 bryozoan faunas, *147*
 bryozoans, 146
 cephalopod genera clustering, *434*
 cluster analysis, *134*
 cluster and ordination tests, 204, 205–207
 number of occurrences, *432*
 Ordovician distribution, 68–69
 Ordovician graptolite biogeography, 419
 time slices, 130–133, *133*, *134*
Volchovia mobilis, *185*
- Darwin, Charles, 26
- Demosponges diversity and palaeobiogeography, 89–90
- dendrogram, 348
- descriptive biogeography, 30
- Deunffia*
 biofacies, 377–378
 distribution, *377*
- Devonian, 365
 graptolite biogeography, 423
 phytoplankton biogeography, 384
- Diaphanometopidae, *307*
- Diaphanometopidae Jaanusson in Moore, 305–306
- Didymaulichnus miettensis* Young, 47
- Didymaulichnus tirasensis*, 47
- Dimeropygidae, *314*
- Dimeropygidae Hupe, 314
- Dionididae, *320*
- Dionididae Gurich, 320
- diplograptid, 423
- discosorids, 440
- disjunct distribution, 31
- dispersed spore fossil record, 461
- distributional dynamics Cambrian clades, 35–40
- diversification of vertebrates in Silurian, 452–455
- diversity curve, *128*
- Dokimocephalidae, *323*
- Dokimocephalidae Kobayashi, 324
- Domasia*
 biofacies, 377–378
 distribution, *377*
- Early Cambrian
 archaeocyathan migration paths, *63*
 Lower Palaeozoic terrane, *10*, *13*, *14*
 palaeogeographical reconstruction, *63*
 phytoplankton, 366–367
- Early Devonian, 6
 Lower Palaeozoic terrane, *20*, *21*
 phytoplankton, 379–380
- Early Early Devonian rostroconchia, 251–253
- Early Late Ordovician ostracod, *350*
- Early Mid Ordovician
 palaeocontinent reconstruction map, *418*
- Early Ordovician
 craniides and *Pseudocrania* geographical distribution, *120*
 Llangynog Inlier, 225–226
 Lower Palaeozoic terrane, *16*
 Montagne Noire, 225
 Montagne Noire and South Wales, 225
- Early Palaeozoic
 biogeography and palaeogeography, 1–2
 biogeography of bryozoans, 145–154
 Conocardiida, *252*
 Icehouse climate, 399–404
 new global palaeogeographical reconstructions, 5–23
 palaeogeographical reconstructions, 14–22
 polychaetes diversification, 266–267
 Rostroconchia, 243–260
 vertebrates palaeobiogeography, 450–456
- Early Palaeozoic vertebrates palaeobiogeography, 450–456
 adaptive radiation of vertebrates in Ordovician, 451–452
 diversification of vertebrates in Silurian, 452–455
 palaeobiological remarks and implications for biostratigraphy, 455
 possible vertebrates of Cambrian, 450–451
- Early Sandbian
 chitinozoan localities, *401*
 Graptolite localities, *401*
- Early Silurian
 Lower Palaeozoic terrane, *19*
 Rostroconchia, 249
- Early to Mid Ordovician, 367–368
 acritarch provincialism, 373–374
 craniiform brachiopods, 121
 provinces, 374
- Early to Mid Palaeozoic
 marine phytoplankton biogeography, 365–389
 Silurian, 376–378
- Eastern Avalonia, 214
- Eastern Gondwana cephalopods, 435
- Eastern Kazakhstania terranes, trilobite, 284

- Eastern Mediterranean region, 278
 East Gondwana palaeogeographical units, 278–281
 echinoderm. *See also* Cambrian echinoderms
 alpha diversity, 166
 Baltica, West Gondwana and Laurentia, 191
 Cambrian, 160, 161
 Cambrian Stage 5, 165
 diversity and Cambrian fluctuations, 166
 first occurrences, 165
 nature of substrates, 174
 ossicles, 164–165
 palaeobiogeographical patterns, 165–166
 plates, 164, 165
 Echinozoans, 185–187
 Echinozoans, 185–187
 Cyclocystoidea, 187
 Echinozoa, 185–186
 Holothuroidea, 186–187
 Ophiocystoidea, 187
 palaeobiogeography of Ordovician echinoderms, 185–187
 Edgewood Province, 107–109, 108
 characteristic corals, 107–108
 coral biogeography in Late Ordovician (Cincinnatian) of Laurentia, 107–109
 coral genera, **101**
 diversity, 108–109
 duration and location, 107
 origination, 108
 termination, 109
 edrioasteroids, 160
 Cambrian echinoderms, 159–161
 Dapingian Stage, 187
 Furongian, 161
 Ordovician echinoderms palaeobiogeography, 187–189
 EiE. *See* Ubiquity hypothesis (EiE)
 Ellesmerocerida, 442
 Encrinuridae, 308
 Encrinuridae Angelin, 307–308
 endemicity, 30, **31**
 Cambrian trilobite palaeobiogeography, 273–290
 clades Laurentia, 166
 pandemic and cosmopolitan distribution, 30–31
 endocerids, 435
 Eocrinoids
 Argentina, 180
 Bolivia, 180
 Cambrian echinoderms, 161–162
 Cambrian Series 3, 162
 Czech Republic, 180
 France, 180
 Furongian, 167
 Morocco, 180
 Spain, 180
 epipelagic species, 362
 eukaryotic taxa, 35
 Eulomidae, 323
 Eulomidae Kobayashi, 323–324
 eunicidan polychaetes, 266
 Euramerican, 378
 Europe ostracod faunas, 356
 extant polychaetes geographical distribution, 270
 Famantina Arc, 19
 Famatina System Argentina, 277
 Famennian acritarchs, 383
 Farewell Terrane, 7, 205
 palaeogeographical reconstructions, 10
 Far Russian East cephalopods, 437
 Fengshan Formation, 441
 Floian, 118
 bryozoans, 146
 cluster analysis, 132
 cluster and ordination tests, 204
 number of occurrences, 432
 Ordovician graptolite biogeography, 417–418
 piloceratids, 435
 stylophorans, 189
 time slices, 129–130, 131
 fold belt, 67
Foliomena fauna, 133
Fordilla, 221
 Fortunian, 54
 Lower Palaeozoic terrane, 10
 France eocrinoids, 180
 Frasnian acritarchs, 383
 Frasnian microfloras, 378
Fritaspis Assemblage, 50
 Furongian
 blastozoan assemblages, 162
 edrioasteroids, 161
 eocrinoids and stylophorans, 167
 Lower Palaeozoic terrane, 15
 oldest scolecodonts, 267
 PAE [Parsimony Analysis of Endemicity] Analysis, 289
 solutans, 163
 stylophorans, 164
 trilobite assemblages, 287
 trilobite genera, 288
 Gamachian, 107
 gastropods, monoplacophorans and mimospirids (GMM), 199–217
 Alaskan areas, 213–214
 Argentinean Precordillera, 216
 Australia, 215
 Baltica and Podolia, 214–215
 cluster and ordination tests, 203–213
 Eastern Avalonia, 214
 gastropod communities, 200–201
 Kazakhstan, 216
 Laurentian areas, 213
 North China, 216
 Ordovician diversification, 222
 Perunica, Morocco and Ibero-Armorica, 215
 Scotland, 214
 Siberia, 216
 Tarim, 215–216
 Geological Time Scale of Gradstein, 470
Gerronodictyon, 74
 ghost lineages, 91–92
 global Cambrian trilobite, 273–290
 global distributional dynamics, 35–40
 global palaeogeographical distribution, 61
 global palaeogeographical reconstructions for Early Palaeozoic, 5–23
 Glyptarcoids Ordovician distributions, 224
 gnathostomes, 453
 Gondwana, 2
 Cambrian Epoch 3, 286
 cluster, 205, 207
 continental blocks and terranes, 281
 ostracods biogeographical patterns, 345
 palaeobiogeographical patterns, 165
 palaeogeographical reconstructions, 6–7
 phylogenetically diverse, 141
 poleward drift, 287
 Silurian, 230–232
 Gorstian cluster and ordination tests, 211–212, 212–213
gracilis interval ostracod genera, **340–343**
 ostracod genera, 344
 ostracods biogeographical patterns, 348
 graptolite, 355
 palaeoecology model, 416
 zone, 345
 zooplankton maps, 399–400
 graptolite palaeobiogeography, 355, 415–425
 biogeographical patterns in graptolite evolution examples, 424
 biogeography revisited, 417
 Devonian graptolite biogeography, 423
 historical biogeography, 423–424
 Ordovician biogeography, 417–420
 palaeobiogeographical models historical development, 415–417

- graptolite palaeobiogeography (*Continued*)
 palaeoecological historical development, 415–417
 Silurian biogeography, 420–423
- Great Ordovician Biodiversification Event (GOBE), 1, 112, 129, 133, 136, 138, 139, 267
- Grewingkia*, 105
Grewingkia canadensis, 106
- Hadoprion cervicornis*, 270
- Hamra Formation, 153
- Harpetidae, 324
- Harpididae, 326
- Harpididae Whittington, 326–327
- hazeliids, 83
- Helicoplacoidea Cambrian echinoderms, 159
- Heminectere*, 82
- Heteroconchs *Lyrodesma poststriatum*, 229–230
 Baltica, 229
 Britain, 229–230
 Kazakhstan, 229
 Ordovician, 229–230
 Siberia, 229
- Heterostracans, 453
- Hexactinellids, 88–89
- hilate spores, 464
- Himalayan margin palaeogeographical units, 280
- Himalayan peri-Gondwanan terranes, 11
- Hirnantian, 345, 375
 bryozoan faunas, 149
 bryozoans, 149
 cluster analysis, 140
 cluster and ordination tests, 208–209
 glaciation, 20
 Ordovician distribution, 73
 Ordovician graptolite biogeography, 420
 ostracod genera, 344
 stage distribution of blastozoans, 183
 time slab ostracods biogeographical patterns, 348–349
 time slices, 136, 140
 zooplankton provinces definitions, 402
- Hirnantian interval, 338
 echinoderm groups, 191
 ostracod genera, 346–347
- Holothuroidea Echinozoans, 186–187
- Holotrachelidae, 315
- Holotrachelidae Warburg, 314–315
- Homalonotidae, 311
- Homalonotidae Chapman, 310
- Homerian cluster and ordination tests, 211–212
- Huanghe type, 436
- Hungaiidae, 326
- Hungaiidae Raymond, 327–328
- Hupeolenus* Zone, 50
- hypercalcified, 67
- hypercalcified taxa, 93
- Hystericuridae, 315
- Hystericuridae Hupe, 315
- Iapetus Ocean islands
 palaeogeographical reconstructions, 10
- Ibero-Armorica
 GMM [Gastropods, monoplacophorans and mimospirids], 215
- ichnospecies, 46
- Illaenidae, 300
- Illaenidae Hawle & Corda, 299
- Illionia*, 233
- Indian craton palaeogeographical units, 280
- Indochina
 palaeogeographical reconstructions, 8–9
- Injetocera, 437
- in situ spores, 463
- International Code of Area Nomenclature, 32
- International Code of Botanical Nomenclature, 27
- International Code of Zoological Nomenclature, 28
- International Subcommittee on Cambrian Stratigraphy (ISCS), 273
- Iran palaeogeographical units, 278–279
- Ischyrinioida, 246
- island arcs, 10
- Isocolidae, 327
- Isocolidae Angelin, 328
- Isorophid edrioasteroids, 187
- Italy palaeogeographical reconstructions, 12–13
- Ityophoridae, 327
- Ityophoridae Warburg, 328
- Jaccard's similarity coefficient, 62
- Junggar Terrane palaeogeographical reconstructions, 11
- Kara
 palaeogeographical reconstructions, 11
 palaeogeographical units, 276–277
 Silurian, 232–234
- Karatau–Naryn trilobite, 284
- Katian
 bryozoan faunas, 148
 cluster and ordination tests, 208
 Lower Palaeozoic terrane, 18
 Ordovician, 374–375
 Ordovician distribution, 70–72
 Ordovician graptolite biogeography, 419–420
 palaeocontinent reconstruction map, 420
 Radiolarian assemblage, 411
 scolecodonts, 268
 stylophorans, 191
- Kazakh
 diversities, 134
 strophomenides, 140
 terranes palaeogeographical reconstructions, 11
- Kazakhstan, 205, 229
 GMM [Gastropods, monoplacophorans and mimospirids], 216
 island arcs and microplates, 279–280
 ostracods biogeographical patterns, 345
 palaeogeographical units, 279–280
 Silurian, 235
 variety of genera, 205
- Klamath terrane, 139
- Kolyma, 276
- Kolymar terrane palaeogeographical reconstructions, 11
- Labechia*, 70
- Labechiella*, 70
- labechiids, 68
- Lady Burn Starfish Bed, 174
- land plants palaeophytogeography, 461–473
- Late Cambrian
 acritarch localities, 369
 global reconstruction, 21
 Lower Palaeozoic terrane, 15
- Late Devonian
 palaeogeography, 383
 phytoplankton, 382–384
- Late Early–early Middle Ordovician rostroconchia, 248
- Late Ludlow–early Přídolí, 359–360
- Late Ordovician
 cephalopod palaeogeography and diversity, 443
 climate, 402–404
 craniiform brachiopods, 121–122
 faunas, 227, 228
 linguliform brachiopods taxa, 120–121
 Lower Palaeozoic terrane, 18
 North America, 228
 Ordovician, 227, 374–376
 palaeocontinent reconstruction map, 420
 rostroconchia, 248–249
 South America, 227–228
 stromatoporoid genera, 70, 71
 stromatoporoids, 69
 zooplankton maps, 399–404
 zooplankton provinces, 401–402
- Late Silurian, 20
 acritarch palaeogeographical reconstruction, 379

- Late Wenlock-early Ludlow, 358–359
- Laurentia
- biogeographical divisions corals, 98
 - brachiopod faunas, 134
 - Ctenocystoids, 163
 - echinoderm assemblage, 191
 - endemic clades, 166
 - Ordovician crinoids, 185
 - ostracods biogeographical patterns, 344–345
 - palaeobiogeographical patterns, 165
 - palaeogeographical reconstructions, 7–8
 - palaeogeographical units, 274–275
 - paracrinoïd, 181
 - paulinitids, 268
 - percentage of shared taxa, **436**
 - plectorthoids, 139
 - rhynchonellide fauna, 136
 - Silurian, 232–234
 - Subrealm, 452
- Laurussia, 8
- Leclerc, Georges-Louis, 26
- Leiostegiidae, 301
- Leiostegiidae Bradley, 300–301
- Lichakephalidae, 302
- Lichakephalidae Tripp, 302–303
- Lichidae, 304
- Lichidae Hawle & Corda, 303–304
- Lichida Moore, 301–304
- Family Lichakephalidae Tripp, 302–303
 - Family Lichidae Hawle & Corda, 303–304
 - synopsis of Ordovician trilobite distribution and diversity, 301–304
- limestone biofacies, cephalopods, 231
- linguliform, 122
- linguliform brachiopods, 117–121
- biogeography of Ordovician, 117–121, 117–124
 - cluster analysis, 120
 - geographical distribution, 118, 119
 - Late Ordovician (Sandbian to Hirnantian), 120–121
 - Mid-Ordovician (Dapingian–Darriwilian), 119–120
- Linnaeus, Carolus, 26
- lithistid, 93
- lithospheric plates, 29
- littoral zone tropical bivalves, 233
- liverwort basal, 463
- Llandeilo, 345
- Llandovery, 472
- bryozoan faunas, 150
 - bryozoans, 150–151
 - Lower Palaeozoic terrane, 19
 - Rostroconchia, 249
 - Silurian distribution, 73–74
 - stromatoporoid genera geographical distribution, 74
 - testing late Wenlock-Prídolí palaeogeography, 360–361
- Llandovery Epoch, 83, 87
- non-stromatoporoid sponges, 88
 - sponge families, **90**
- Llangynog Inlier, 225–226
- Ludlovian bryozoan faunas, 152
- Lophiostroma*, 76
- Lower and Middle Ordovician radiolarian, 409
- Lower Katian
- cluster analysis, 137
 - time slices, 134–135, 137
- Lower Ordovician radiolarian, 407–408
- distribution, 409
- Lower Palaeozoic
- Ordovician, 222–230
 - outcrops, 8
 - palaeobiogeography of *Bivalvia*, 221–236
 - palaeogeographical reconstructions, 6–8
 - Silurian, 230–235
 - terrane boundaries, 6, 7, 9
- Lower Palaeozoic terrane
- Cambrian–Ordovician boundary, 15
 - Early Cambrian, 13, 14
 - Early Cambrian, Fortunian, 10
 - Early Devonian, Emsian, 21
 - Early Devonian, Pragian, 20
 - Early Ordovician, Tremadocian, 16
 - Early Silurian, Llandovery, Rhuddanian, 19
 - Early Silurian, Llandovery, Telychian, 19
 - Late Cambrian, Furongian, 15
 - Late Ordovician, Katian, 18
 - Late Ordovician, Sandbian, 18
 - Late Silurian, Ludlow, Ludfordian, 20
 - Middle Cambrian, 14
 - Middle Ordovician, 17
 - Middle Ordovician, Floian, 16
- Ludfordian, 20
- cluster and ordination tests, 212–213
- Ludlow, 20
- bryozoans, 152–154
 - Silurian distribution, 74–77
 - stromatoporoid genera geographical distribution, 76
- Ludlow Epoch, 83
- non-stromatoporoid sponges, 89
- Ludlow-Prídolí
- graptolites, 421
 - sponge families, **92**
- Ludlow Series, 378
- Lyell, Charles, 26
- Mahalospongiidae, 90
- Major Early Palaeozoic events, 14–22
- Cambrian, 14–16
 - Ordovician, 16–20
 - Silurian, 22
- Malongulli Formation, 411
- Malopolska, 347
- marine phytoplankton, 365–389
- Mediterranean Province, 368–369
- megafossil record, 461
- Megagraption*, 50
- Megalomoidea, 233
- Mesaktoceras* occurrences, 440
- mesopelagic species, 362
- metric multidimensional scaling (MDS), 471
- plot, 472
- Mexican terranes palaeogeographical reconstructions, 11
- microcontinents, **28**
- microplates, **28**
- palaeogeography in tectonics, 28–29
- Mid and late Cambrian phytoplankton, 366–367
- Mid Devonian
- palaeogeography, 381
 - phytoplankton, 380–382
- Middle Cambrian, 14
- Middle Eastern terranes palaeogeographical reconstructions, 11
- Mid Ordovician
- Australia, 226
 - faunas, 226–227, 226
 - linguliform brachiopods taxa, 119–120
 - Lower Palaeozoic terrane, 16, 17
 - Ordovician cephalopod palaeogeography and diversity, 442–443
 - Radiolarian, 409–410
 - Spain, 226
 - stromatoporoid genera, 68
 - Wales, 226–227
 - West Yunnan, 227
- Mid Silurian
- distribution Megalomoidea, 233
 - distribution Praecardiina, 232
 - Silurian, 376–378
- mimospirids. *See* Gastropods, monoplacophorans and mimospirids (GMM)
- Missisquoiidae, 302
- Missisquoiidae Hupe, 301
- mixed-layer organisms, 351
- moderate endemism model, 389
- Moldanubia palaeogeographical reconstructions, 12–13
- Mollusca. *See* Rostroconchia

- Mongolian terranes palaeogeographical reconstructions, 11
monographtid, 424
Monomorphichnus, 52
monoplacophorans. *See* Gastropods, monoplacophorans and mimospirids (GMM)
Montagne Noire, 225
 early Ordovician faunas, 225
 faunas, 225
Moroccan Atlas Cambrian strata, 278
Morocco
 eocrinoids, 180
 GMM [Gastropods, monoplacophorans and mimospirids], 215
myodocopes, 356
- nappe structures, 22
nearshore cluster, 351
Nemagraptus gracilis, 419
Neoproterozoic basement, 10
Neoproterozoic-Cambrian succession, 280
Neoveryhachium carminae
 biofacies Silurian, 376–377
 distribution, 377
Nereites, 50
New Zealand palaeogeographical units, 281
Nileidae, 319
Nileidae Angelin, 319
non-stromatoporoid Porifera, 81–93
non-stromatoporoid sponges
 distribution for Llandovery Epoch, 88
 distribution for Ludlow and Přídolí epochs, 89
 distribution for Wenlock Epoch, 89
 Přídolí epochs, 89
 Wenlock Epoch, 89
North America
 Late Ordovician faunas, 228
 terranes palaeogeographical units, 277–278
 testing Silurian palaeogeography, 356
North China
 cephalopods, 436–437
 GMM [Gastropods, monoplacophorans and mimospirids], 216
 palaeogeographical reconstructions, 11–12
 palaeogeographical units, 280–281
Nuculoids Ordovician, 228–229
- oceans biogeographical regions and climate regions, 27
Odontopleurida, 301
Odontopleuridae, 304
Odontopleuridae Burmeister, 304
Odontopleurida Whittington in Moore, 304
 synopsis of Ordovician trilobite distribution and diversity, 304
Odontopleurida Whittington in Moore
 Family Odontopleuridae Burmeister, 304
Oldhamia, 47
 distribution, 49
 Forbes, 47–49
Old Red Sandstone Continent, 454
Olenida Adrain, 321–324
 Family Dokimocephalidae Kobayashi, 324
 Family Eulomidae Kobayashi, 323–324
 Family Olenidae Burmeister, 321–322
 Family Remopleuridae Hawle & Corda, 322–323
 Family Sarkiidae Hupe, 324
 synopsis of Ordovician trilobite distribution and diversity, 321–324
Olenidae, 322
Olenidae Burmeister, 321–322
Omolon Terrane palaeogeographical reconstructions, 11
oncocerids, 440
Ophiocistiioidea Echinozoans, 187
Ophiuroids Asterozoans, 178
ordination tests, 203–213
Ordovician, 222–230, 367–376
 Baltic Province, 369
 biogeography of early to mid Palaeozoic phytoplankton, 367–376
 biogeography of GMM [Gastropods, monoplacophorans and mimospirids], 199–217
 bivalve distribution, 223
 blastozoans, 183
 blastozoans morphology, 179, 182
 cephalopod morphologies, 430
 cephalopod palaeogeography and diversity, 429–444
 cephalopods diversity trends, 436
 diploporans palaeobiogeographical distribution, 180
 dispersed spore diversity curve, 465
 distributions, 224–225
 diversification of gastropods and bivalves, 222
 duration of Early to Mid Ordovician Provinces, 374
 early Ordovician faunas, 225–226
 Early to Mid Ordovician, 367–368
 Early to Mid Ordovician acritarch provincialism, 373–374
 faunal composition, 225
 gastropod biocommunities, 201
 Heteroconchs *Lyrodesma poststriatum*, 229–230
 Hirnantian, 375–376
 holothurian species, 186
 land plants palaeophytogeography, 461–473
 Late Ordovician, 374–376
 Late Ordovician faunas, 227
 linguliform and craniiform brachiopods biogeography, 117–124
 Lower Palaeozoic palaeobiogeography of Bivalvia, 222–230
 Major Early Palaeozoic events, 16–20
 mass extinction, 97
 Mediterranean Province to Perigondwana Province, 368–369
 mid-Ordovician faunas, 226–227
 NMDS diagram, 210
 non-stromatoporoid Porifera, 81–93, 86–91
 Nuculoids, 228–229
 palaeogeographical regions of cephalopods, 432–438
 polychaete diversity and biogeography, 265–271
 polychaetes global genus-level diversity, 268
 Radiolarian palaeogeographical distribution, 407–412
 rhombiferans palaeobiogeographical distribution, 181
 rostronch species, 254–260
 Sandbian and Katian, 374–375
 scolecodont-bearing polychaete genera, 266
 South China, Perigondwana, or Baltica, 369–370
 sponge palaeobiogeography Porifera, 84–86
 sponges distribution, 85
 trilobite, 297–329
 trilobite distribution and diversity, 297–329
 trilobite families stratigraphical ranges, 298
 vertebrates adaptive radiation, 451–452
 zooplankton, 399–401
 zooplankton maps and climate of the Early Palaeozoic Icehouse, 399–404
Ordovician asterozoans
 morphology, 177
 palaeobiogeographical distribution, 178
Ordovician biodiversity, biogeography and phylogeography
 biogeographical provinces, 128–129
 brachiopod biodiversity and phylogeography during Ordovician, 136–142
 rhynchonelliform brachiopods, 127–142
 time slices, 129–136
Ordovician crinoids
 Laurentia, 185
 morphology, 184
 palaeobiogeographical distribution, 184
Ordovician distribution, 67–73, 224–225
 Afghanodesmatids, 224
 Ambonychids, 225
 biogeography of the Ordovician and Silurian Stromatoporoidea, 67–73
 Cyrtodontids, 224–225
 Darrwilian, 68–69
 Glyptarcoids, 224
 Hirnantian, 73
 Redoniids, 224
 Sandbian, 69–70
Ordovician echinoderms
 Baltica, 174, 186
 disparity patterns, 174
 evolution, 176
 global and regional biodiversity patterns, 174

- palaeobiogeographical analysis, 176
 record, 177
 Ordovician echinoderms palaeobiogeography, 173–192
 Asterozoans, 177–179
 Blastozoans, 179–184
 Crinoids, 184–185
 database, 175
 Echinozoans, 185–187
 Edrioasteroids, 187–189
 palaeobiogeographical maps, 177
 quality of dataset and possible biases, 175–177
 Stylophorans, 189–192
 Ordovician echinozoans
 palaeobiogeographical distribution, 186
 Ordovician edrioasteroids
 morphology, 188
 palaeobiogeographical distribution, 188
 Ordovician eocrinoids
 palaeobiogeographical distribution, 182
 Ordovician facies zones
 Baltoscandia, 338
 Ordovician faunas
 palaeobiogeography, 268–269
 polychaete diversity and biogeography, 268–269
 Ordovician genera
 distribution, **205**
 Ordovician generic brachiopod diversity and phylogeographical
 distribution, 141
 Ordovician graptolite biogeography, 417–420
 Dapingian, 418–419
 Darriwilian, 419
 Floian, 417–418
 Hirnantian, 420
 Katian, 419–420
 Tremadocian, 417
 Ordovician ostracods
 biogeographical patterns, 337–351
 biogeography, 350
 cluster analysis, 350
 correspondence analysis, 350
 Ordovician stylophorans
 morphology, 189
 palaeobiogeographical distribution, 190
 Orthida Klamath terrane, 139
 Ossa-Morena trilobite, 283
 Ostracoda, 337
 ostracoderms, 454
 ostracod genera
 gracilis interval, **340–343**
 Hirnantian, 344
 Hirnantian interval, **346–347**
 ostracods biogeographical patterns, 337–351
 biogeography, 349–351
 data analysis, 347–348
 database, 347
 Gondwana, 345
 gracilis time slab, 348
 Hirnantian time slab, 348–349
 Kazakhstan, 345
 Laurentia, 344–345
 Ordovician, 337–351
 palaeogeography, 345–347
 Peri-Gondwana, 345
 selection of time slabs, 338
 Siberia and adjacent areas, 345
 source data, 338–345
 taxonomic remarks on source data, 345
 ostracod species
 palaeogeographical distribution, 361
 stratigraphical distribution, 357
 Otta terrane, 205
 Ouachita Suture, 11
 overlap zone, 123

 palaeobiogeographers, 26–27
 palaeobiogeographical nomenclature
 proposal for consistent use, 31–32
 ranks, 31–32
 standardization, 32
 palaeobiogeographical patterns, 165–167
 alpha diversity in space and time, 166–167
 biogeographical relationships of echinoderm assemblages,
 165–166
 Cambrian echinoderm diversity and palaeobiogeography,
 165–167
 first appearances in Laurentia and Gondwana, 165
 palaeobiogeography, 25–28, 30–31
 biogeographical entities: biomes and regions, 27
 biogeographical units, 31
 biogeography defined, 25–26
 Cambrian trilobite assessed using parsimony analysis of endemism,
 273–290
 defined, 30
 Early Palaeozoic vertebrates, 450–456
 ecological entities: communities and ecosystems, 27–28
 endemic, pandemic and cosmopolitan distribution, 30–31
 first biogeographers and palaeobiogeographers, 26–27
 nomenclatural rules and problems, 28
 Ordovician and Silurian polychaete diversity and biogeography,
 267–269
 Ordovician echinoderms, 173–192
 Ordovician faunas, 268–269
 provincialism, 31
 Siberian faunas, 269
 terminology, 30
 palaeobiogeography of Bivalvia, 221–236
 palaeoecology, 60–62
 palaeogeographical distribution
 Botomian Stage, 61
 Cambrian echinoderms, 159–164
 Ordovician radiolarian occurrences, 407–412
 pattern of Rostroconchia, 246–253
 Silurian benthic ostracod species, 359
 palaeogeographical positions Cambrian localities, **37**
 palaeogeographical reconstructions
 Annamia (Indochina), 8–9
 and Archaeocyatha migration paths, 63
 Arctic Alaska–Chukotka, 9
 Armorican Terrane Assemblage, 9–10
 Avalonia, 10
 Baltica, 8
 Bruno–Silesia, 12–13
 Cordilleran terranes, 10
 Early Palaeozoic and their generation, 5–23
 Farewell Terrane, 10
 Gondwana, 6–7
 Iapetus Ocean islands, 10
 Italy, 12–13
 Junggar Terrane, 11
 Kara, 11
 Kazakh and Tien Shan terranes, 11
 Kolymar–Omolon Terrane, 11
 Laurentia, 7–8
 major continental blocks in Lower Palaeozoic, 6–8
 Major Early Palaeozoic events, 14–22
 Mexican and Central American terranes, 11
 Middle Eastern and Himalayan peri-Gondwanan terranes, 11
 Moldanubia, 12–13
 Mongolian terranes, 11
 North China (Sinokorea), 11–12
 other terranes, 8–13
 Palaeoaltitudes of localities, **348**
 Pearya, 12
 Peri-Gondwanan terranes now to the south, 12
 Perunica (Bohemia), 12
 Pontides and Taurides, 12
 Precordillera, 12
 Reno–Hercynian Terrane, 12–13
 Saxothuringia, 12–13
 Siberia, 8
 Sibumasu (Shan–Thai), 13
 South China, 13

- palaeogeographical reconstructions (*Continued*)
 Svalbard and adjacent terrane areas, 13
 Tarim, 13
 Tien Shan terrane, 11
- palaeogeographical regions, 432–438
- palaeogeographical units, 274–281
 Afghanistan, 279
 Altai-Sayan area, 276
 Australia–New Zealand–Antarctica, 281
 Baltica, 274
 Central American terranes, 277
 Central Asia, 279
 Eastern Mediterranean region, 278
 East Gondwana, 278–281
 Indian craton and Himalayan margin, 280
 Iran, 278–279
 Kara terrane, 276–277
 Kazakhstani island arcs and microplates, 279–280
 Laurentia, 274–275
 North American terranes, 277–278
 Saxo-Thuringia and Bohemia, 278
 Siberia, 275–277
 Siberian Platform, 276
 South America, 277
 South China and Tarim, 280–281
 Western Mediterranean terranes, 278
 West Gondwana, 277–278
- palaeogeography
 Archaeocyatha in Cambrian biostratigraphy and biogeography, 62–63
 ostracods biogeographical patterns, 345–347
- palaeogeography in tectonics, 28–30
 basic terminology, 28–30
 continent – microcontinent – craton, 29
 impact of plate tectonic theory on palaeobiogeography, 28
 plates and microplates, 28–29
 terrane – terrane assemblage, 29–30
- palaeophytogeography
 dispersed spores, 466–467
 early land plant fossil record, 461–463
 early land plant palaeophytogeography, 466–467, 469–473
 general principles, 467–469
 interpretation of new database/analyses, 473
 megafossils, 466
 nature of early land plant adaptive radiation, 464–466
 Ordovician–Silurian land plants, 461–473
 origin and early diversification of land plants, 463–464
- palaeoplates, 29
- Palaeozoic marine phytoplankton biogeography, 365–389
- Palaeozoic palaeogeographical and palaeobiogeographical nomenclature, 25–32
 basic terminology in palaeo(bio)geography, 30–31
 basic terminology of (palaeo)geography in tectonics, 28–30
 origins and outline of (palaeo)biogeography, 25–28
 palaeobiogeographical nomenclature standardization, 32
 proposal for consistent use of palaeobiogeographical nomenclature, 31–32
- Palaeozoic terrane boundaries, 8
- Paleodictyon*, 51
- Paleodictyon meneghini*, 49–50
- panbiogeography, 27
- pandemic distribution, 31
- pandemism, 31
- Panderiidae, 300
- Panderiidae Bruton, 299
- paracrino Laurentian Province, 181
- Paredriophus elongatus*, 187
- Parsimony Analysis of Endemicity (PAE), 2
 Cambrian Epoch 3, 289
 global Cambrian trilobite palaeobiogeography assessed, 273–290
- parsimony-based analysis, 423
- paulinitids Laurentia, 268
- Pearya palaeogeographical reconstructions, 12
- pelagic ostracod species distribution, 361
- peri-continental terranes, 6
- Perigondwana, 277
 Ordovician, 368–369, 369–370
 ostracods biogeographical patterns, 345
 palaeogeographical reconstructions, 12
 terranes, 7
- Perunica
 cephalopods, 433
 GMM [Gastropods, monoplacophorans and mimospirids], 215
 palaeogeographical reconstructions, 12
- Phacopidae, 306
- Phacopidae Hawle & Corda, 305
- Phacopida Salter, 304–310
 Family Acastidae Delo, 304–305
 Family Bathycheilidae Přibyl, 309
 Family Bavarillidae Sdzuu, 309
 Family Calymenidae Swinerton, 309–310
 Family Cheiruridae Hawle & Corda, 306
 Family Dalmanitidae Vogdes, 305
 Family Diaphanometopidae Jaanusson in Moore, 305–306
 Family Encrinuridae Angelim, 307–308
 Family Homalonotidae Chapman, 310
 Family Phacopidae Hawle & Corda, 305
 Family Pharostomatidae Hupe, 310–311
 Family Pliomeridae Raymond, 308–309
 Family Prosopiscidae Fortey & Shergold, 306
 Family Pterygometopidae Reed, 305
 Suborder Calymenina Swinerton, 309–310
 Suborder Cheirurina Harrington & Leanza, 306–309
 Suborder Phacopina Salter, 304–306
 synopsis of Ordovician trilobite distribution and diversity, 304–310
- Phacopina Salter, 304–306
- Pharostomatidae, 311
- Pharostomatidae Hupe, 310–311
- phosphatic microremains, 450
- phylogeography of Ordovician, 127–142
- phytoplankton, 365–389
 biogeography of early to mid Palaeozoic, 365–389
 Cambrian phytoplankton, 366–367
 changes in biogeographical ranges through time, 387–388
 continental margin distributions, 387
 cosmopolitanism, endemism and community breakdown, 388
 distal v. proximal (oceanic v. cratonic) distribution patterns, 385–386
 Early Cambrian phytoplankton, 366–367
 Early Devonian, 379–380
 Late Devonian, 382–384
 latitude v. other controls, 384–385
 Mid and late Cambrian phytoplankton, 366–367
 Mid Devonian, 380–382
 Ordovician, 367–376
 Silurian, 376–378
 synopsis of Devonian phytoplankton biogeography, 384
- plates tectonic theory, 28
 palaeobiogeography, 28
 palaeogeography, 28–29
- plectorthoids Laurentia, 139
- Pliocene Research, Interpretation and Synoptic Mapping (PRISM), 399
- Pliomeridae, 309
- Pliomeridae Raymond, 308–309
- Plumatalinia*, 71, 73
- podocope, 362
- Podolia, 214–215
- Pojetaia*, 221
- polychaete annelid worms, 265
- polychaete diversity and biogeography
 Early Palaeozoic polychaetes and their diversification, 266–267
 Ordovician, 265–271
 Ordovician and Silurian, 265–271
 Ordovician faunas, 268–269
 palaeobiogeographical units and distribution patterns, 269–270
 palaeobiogeography, 267–269
 Siberian faunas, 269
- Pontides palaeogeographical reconstructions, 12
- Porifera, 59, 81–93
 Cambrian, Ordovician and Silurian, 81–93
 Cambrian sponge palaeobiogeography, 83–84
 ghost lineages, 91–92

- modern sponges distribution patterns, 81–82
 non-stromatoporoid, 81–93
 Ordovician sponge palaeobiogeography, 84–86
 Silurian palaeobiogeography, 92–93
 Silurian sponge diversity and palaeobiogeography, 86–91
 taphonomic influence on perceived sponge distributions, 92
 taxonomic and taphonomic problems, 82
- Pradesura*, 178
 Praecardiidina Mid-Silurian distribution, 232
 praecardiidid bivalves, 231
 Pragian Lower Palaeozoic terrane, 20
 Precambrian cores, 20
 Precordillera
 cephalopods, 438
 palaeogeographical reconstructions, 12
 pre-Hirnantian Ashgill, 135
 pre-trilobitic *Rusophycus* Hall, 50–52
 priapulid-like producer, 47
 Přídolí, 77
 bryozoan faunas, 153
 bryozoans, 154
 Přídolí Epoch, 83
 non-stromatoporoid sponges, 89
 Přídolí MDS, 473
 Přídolí stromatoporoid genera, 77
 Proetidae, 312
 Proetidae Salter, 311–312
 Proetida Fortey & Owens, 311–312
 Family Proetidae Salter, 311–312
 Family Tropicocoryphidae Přibyl, 312
 synopsis of Ordovician trilobite distribution and diversity, 311–312
 prokaryotic taxa, 35
Propora, 110
 Prosopiscidae, 307
 Prosopiscidae Fortey & Shergold, 306
Protaster, 178
 protocycloceratids, 435
Protolenus Zone, 49
 Protomonaxonids, 90
Protospongia, 82, 88
 provincialism, 77
Pseudocrania, 120
Pteraspis, 455
 Pterygometopidae, 306
 Pterygometopidae Reed, 305
- Queenston Delta, 107
- radiocyathan genera cluster analysis, 62
 Radiolarian, 407–412
 distribution of assemblages, 407–411
 Lower Ordovician assemblages, 407–408
 Middle Ordovician assemblages, 409–410
 palaeoenvironmental influence on faunal composition, 411–412
 palaeogeographical distribution of Ordovician, 407–412
 patterns, significance and limitations, 407–412
 quality of current data, 411–412
 stratigraphical accuracy, 411
 taphonomic biases, 412
 taphonomic matters, 412
 Upper Ordovician assemblages, 410–411
 Raphiophoridae, 321
 Raphiophoridae Angelin, 320
 Raymondinidae, 328
 Raymondinidae Clark, 329
 Redoniids Ordovician distributions, 224
 Red River-Stony Mountain Province, 99–104
 characteristic corals, 102
 coral biogeography in Late Ordovician (Cincinnatian) of Laurentia, 99–104
 coral genera, **100**
 diversity, 102–104
 duration and location, 99–100
 origination, 102
 termination, 104
 reefs biosedimentology and lithology, 62
 Remopleurididae, 322
 Remopleurididae Hawle & Corda, 322–323
 representative palaeogeographical units, 274–281
 Revised Paleozoic APW paths, 5
Rhegmaphyllum, 110
 Rheic Ocean, 384
 Rheno–Hercynian Terrane palaeogeographical reconstructions, 12–13
 Rhombifera Cambrian echinoderms, 162
 Rhuddanian
 cluster and ordination tests, 208–209
 Rhynchonellida, 133
 fauna, 136
Rhytiprion magnus, 270
 Ribeirioida, 243, 244, 246
 Richmondian, 99
 Richmondian Stage, 105
 Richmond Province, 105–107
 characteristic corals, 105
 coral biogeography in Late Ordovician (Cincinnatian) of Laurentia, 105–107
 coral genera, **101**
 diversity, 106
 duration and location, 105
 origination, 106
 termination, 107
 Rorringtoniidae, 316
 Rorringtoniidae Owens in Owens & Hamann, 315–316
 Rostroconchia, 2, 243–260
 biogeographical distribution patterns in Early Palaeozoic, 243–260
 biological factors, 245
 distribution, 247, 248, 249, 250
 distribution map, 247, 250, 251
 Early Early Devonian (Lochkovian–early Pragian), 251–253
 Early Early Ordovician (Tremadocian), 247–248
 Early Silurian (Llandovery), 249
 Late Early–early Late Silurian (Wenlock–Ludlow), 249–251
 Late Early–early Middle Ordovician (Floian–Dapingian), 248
 Late Ordovician (Sandbian–Hirnantian; Caradoc, Ashgill), 248–249
 Later stages (late Early Devonian–Middle Devonian), 253
 palaeogeographical analysis, 245
 Palaeogeographical distribution pattern, 246–253
 preservation, 244
 taxonomic and systematic basis, 246
 Rugosa, 97, 98
Rusophycus Hall, 51
 testing for palaeogeographical patterns in distribution, 50–52
- Sacabambaspis* artist's reconstruction, 451
Saloceras occurrences, 440
 Sandbian, 17
 biotope, 401
 bryozoan faunas, 147
 bryozoans, 146
 climate belts, 403
 cluster analysis, 136
 cluster and ordination tests, 205–207
 Lower Palaeozoic terrane, 18
 Ordovician, 374–375
 Ordovician distribution, 69–70
 palaeoceanographical boundaries, 402
 stylophoran assemblages, 190
 time slice, 133–134, 135
 zooplankton provinces definitions, 401–402
 Sandbian-early Katian cephalopod clustering, 434
 Sandbian-Katian
 blastozoans, 181
 cluster and ordination tests, 207–208
 number of occurrences, 432
 Sandbian Stage, 69
 Sarkiidae, 324
 Sarkiidae Hupe, 324
 Saxothuringia
 palaeogeographical reconstructions, 12–13
 palaeogeographical units, 278
 Scharysiidae, 316
 Scharysiidae Osmólska, 316

- Sclater, Philip Lutley, 26
- scolecodont-bearing polychaete genera
 Ordovician, 266
 Silurian, 267
- scolecodonts
 South China, 268
 Tarim, 268
- Scotland, 214
- Scutelluidae, 300
- sea urchins. *See* Echinoids
- Selty terrane trilobite, 284
- Sheinwoodian-Homerian, 211
- Shumardiidae, 328
- Shumardiidae Lake, 329
- Shundy Formation of Kazakhstan, 409
- Siberia
 cephalopods, 437
 faunas palaeobiogeography, 269
 GMM [Gastropods, monoplacophorans and mimospirids], 216
 Heteroconchs *Lyrodesma poststriatum*, 229
 ostracods biogeographical patterns, 345
 palaeogeographical reconstructions, 8
 palaeogeographical units, 275–277
 records each geographical area, 211
- Siberian Caucasus Basin, 235
- Siberian Platform palaeogeographical units, 276
- Sibumasu
 cephalopods, 435
 palaeogeographical reconstructions, 13
- Silurian, 199–217, 230–235, 376–378
 biogeography of early to mid Palaeozoic, 376–378
 biogeography of GMM [Gastropods, monoplacophorans and mimospirids], 199–217
 bryozoans, 149–150
 Deunffia-Domasia biofacies, 377–378
 Devonian gastropod communities, 201
 dispersed spore diversity curve, 465
 dispersed spores, 471
 distribution of dispersed spores, 469–470
 diversification of vertebrates, 452–455
 gastropod biocommunities, 201
 Gondwana, 230–232
 graptolite biogeography, 420–423
 Kazakhstan, 235
 land plants palaeophytogeography, 461–473
 Laurentia, Avalonia, Baltica and Kara, 232–234
 Lower Palaeozoic palaeobiogeography of Bivalvia, 230–235
 Major Early Palaeozoic events, 22
 Mid Silurian (late Llandovery-Wenlock), 376–378
Neoveryhachium carminae biofacies, 376–377
 NMDS diagram, 212
 non-stromatoporoid Porifera, 81–93, 86–91
 non-stromatoporoid sponges, 88
 ostracods, 355
 palaeobiogeography Porifera, 92–93
 palaeocontinental reconstructions, 467
 pelagic ostracod species, 361
 polychaete diversity and biogeography, 265–271
 relative endemism, 204
 rostroconch species, 254–260
 scolecodont-bearing polychaete genera, 267
 Siberia, 234–235
 sponge families, 87
 vertebrates biogeography, 454–455
- Silurian benthic ostracod species
 palaeogeographical distribution, 359
 stratigraphical distribution, 357
- Silurian distribution, 73–77
 Llandovery, 73–74
 Ludlow, 74–77
 Přídolí, 77
 Wenlock, 74
- Silurian palaeogeography
 Benthic species, 362–363
 comparators of recent benthic and pelagic ostracods, 362–363
 lifestyles and biogeographical regions of Silurian ostracods, 355
 Pelagic species, 362–363
 testing late Wenlock-Přídolí, 356–360, 360–362
 using 'European' ostracod faunas, 355–363
- Silurian sponge diversity and palaeobiogeography, 86–91
 Calcareans, 91
 Cambrian, Ordovician and Silurian non-stromatoporoid Porifera, 86–91
 Demosponges, 89–90
 Hexactinellids, 88–89
 Porifera, 86–91
 Protomonaxonids, 90
- Sino-Korean block, 281
- Skiagia-bearing assemblage distribution, 367
- Slavidae, 231
- Soluta Cambrian echinoderms, 163
- Somasteroids Asterozoans, 177–178
- South America
 Late Ordovician faunas, 227–228
 palaeogeographical units, 277
- South China
 cephalopods, 437–438
 Ordovician, 369–370
 palaeogeographical reconstructions, 13
 palaeogeographical units, 280–281
 Palaeoplate, 137
 scolecodonts, 268
- South Wales Early Ordovician faunas, 225
- Spain
 eocrinoids, 180
 mid-Ordovician faunas, 226
 species pump, 129, 139
Sphinctozoan demosponges, 90
 spinose siphonretides, 122
 sponge families
 distribution, 82
 distribution for Llandovery Epoch, 90
 distribution for Ludlow-Přídolí epochs, 92
 distribution for Wenlock Epoch, 91
 distribution patterns, 81–82
 diversity and palaeobiogeography, 86–91
 species and genera Silurian Period, 87
- spore
 assemblages, 464
 morphological disparity curve, 465
- Stairway Sandstone, 227
- Stenaster*, 178
- stenohaline organisms, 60
- Streptelasma divaricans*, 106
- strike-slip faulting, 8
- stromatoporoids
 distribution of early Late Ordovician, 69, 70
 distribution of late Mid-Ordovician, 68
 distribution of mid-late Late Ordovician, 70, 71
 recorded occurrences, 73, 75
- strophomenide
 clades explosion, 138
 Kazakh terranes, 140
- Styginidae, 301
- Styginidae Vogdes, 299–300
- stylophora
 Cambrian echinoderms, 163–164
 Floian, 189
 fossil record, 190
 Furongian, 164, 167
 Katian, 191
 Ordovician echinoderms palaeobiogeography, 189–192
 Sandbian, 190
- Sunnaginia imbricata* Zone, 49
- Svalbard and adjacent terrane areas, 13
- Symmetropirion*, 270
- Syringomorpha nilssoni*
 palaeogeographical patterns, 52–53
 stratigraphical distribution, 53
- Tabulata, 97
- Taconic Orogeny, 18

- Taihungshaniidae, 319
 Taimyr Peninsula, 11
 taphonomic bias, 46
 Tarim, 205
 Cambrian strata, 281
 cephalopods, 438
 GMM [Gastropods, monoplacophorans and mimospirids], 215–216
 palaeogeographical reconstructions, 13
 palaeogeographical units, 280–281
 scolecodonts, 268
 Tarphycerida, 432
 tarphycerids, 436
 Taugantelyski Formation, 153
 Taurides
 palaeogeographical reconstructions, 12
 tectonics terminology, 28
 tectonostratigraphic terrane, 30
 Telephinidae, 317
 Telephinidae Marek, 316–317
 Telychian, 19
 Telychian-Sheinwoodian, 210
 terrane, 28
 palaeogeography in tectonics, 29–30
 Terreneuvian Series, 366
 Tien Shan terrane, 11
 palaeogeographical reconstructions, 11
 time slabs, 338
 time slices, 129–136
 biodiversity, biogeography and phylogeography of Ordovician, 129–136
 Dapingian–Darriwilian, 130–133
 Floian, 129–130
 Hirnantian, 136
 Lower Katian, 134–135
 Sandbian, 133–134
 Tremadocian, 129
 Upper Katian, 135–136
 Tommotian Stage, 59
 global palaeogeographical distribution, 61
 torynelasmatids, 119
 trace fossils. *See* Cambrian trace fossils
 transgressive-regressive cycles, 111
 Tremadocian, 337, 436
 bryozoans, 146
 Lower Palaeozoic terrane, 16
 Ordovician bivalves, 223
 Ordovician graptolite biogeography, 417
 Tremadocian Age, 118
 Tremadocian cephalopods, 439
 Tremadocian-Floian, 203–204
 Tremadocian Stage, 137
 Tremadocian stylophorans, 189
 Tremadocian time slice, 130
 cluster analysis, 130
 Tremadoc Series crinoids, 184
 trilete spores, 464
 trilobite, 16. *See also* Cambrian trilobite
 Alai terrane, 283–284
 archaeocyaths, 60
 Avalonia, 282–283
 based biogeographical information support terrane distinctions, 281–284
 Cuyania, 283
 distribution and diversity, 297–329
 Family Acrocephalitidae Hupe, 325–326
 Family Catillicephalidae Raymond, 326
 Family Harpididae Whittington, 326–327
 Family Hungaiidae Raymond, 327–328
 Family Isocolidae Angelin, 328
 Family Ityophoridae Warburg, 328
 family-level histories, 299–329
 Family Raymondinidae Clark, 329
 Family Shumardiidae Lake, 329
 Furongian, 287
 global Cambrian trilobite palaeobiogeography, 281–284
 Kazakhstanian island arcs and microplates, 284
 Order Asaphida Salter, 317–321
 Order Aulacopleurida Adrain, 312–317
 Order Corynexochida Kobayashi, 299–301
 Order Harpida Whittington, 324–325
 Order Lichida Moore, 301–304
 Order Odontopleurida Whittington in Moore, 304
 Order Olenida Adrain, 321–324
 Order Phacopida Salter, 304–310
 Order Proetida Fortey & Owens, 311–312
 Order uncertain, 325–329
 Ordovician, 297–329
 Ossa-Morena, 283
 summed diversity, sampling and endemism, 297–299
 trimerellides, 121
 trimerellide taxa geographical distribution, 121
 Trinucleidae, 321
 Trinucleidae Hawle & Corda, 319–321
 Trocholites, 433, 438
 Tropidocoryphidae, 312
 Tropidocoryphidae Přibyl, 312
 Tuaringia, 221
 Tuva, 276
 ubiquity hypothesis (EiE), 389
 unweighted pair-group method using arithmetic averages (UPGMA), 388
 Upper Katian
 cluster analysis, 139
 time slices, 135–136, 138
 Upper Ordovician assemblages, 410–411
 Upper Palaeozoic outcrops, 8
 vertebrates in Silurian
 biogeography of Silurian vertebrates, 454–455
 climatic context, 455
 diversification, 452–455
 fossil record, 452–454
 palaeobiogeography of Early Palaeozoic vertebrates, 452–455
 vertebrates palaeobiogeography, 450–456
 Volchovia mobilis, 185
 von Humboldt, Alexander, 26
 Wales mid-Ordovician faunas, 226–227
 Wegener, Alfred, 26
 Wenlock
 bryozoan faunas, 151
 bryozoans, 151–152
 Silurian distribution, 74
 Wenlock Epoch, 83, 89
 non-stromatoporoid sponges, 89
 sponge families, 91
 Wenlock-Ludlow/Přídolí
 testing palaeogeography, 361–362
 Wenlock-Přídolí
 Late Ludlow-early Přídolí, 359–360
 Late Wenlock-early Ludlow, 358–359
 Llandovery, 360–361
 palaeogeography, myodocope ostracods, 360–362
 palaeogeography, palaeocopes and podocopes, 356–360
 testing Silurian palaeogeography using ‘European’ ostracod faunas, 360–362
 Wenlock-Ludlow/Přídolí, 361–362
 Wenlock stromatoporoid genera geographical distribution, 75
 Western Mediterranean terranes, 278
 West Gondwana
 Avalon, 282
 cephalopods, 438
 echinoderm assemblage, 191
 palaeogeographical units, 277–278
 West Yunnan mid-Ordovician faunas, 227
 Wildenow, Karl, 26
 Wutinoceras occurrences, 440
 xiphograptid, 424
 York terrane, 207

zooplankton maps

- BugPlates palaeogeographical reconstruction, 401
- chitinozoans, 400–401
- and climate of the Early Palaeozoic Icehouse, 399–404
- graptolites, 399–400
- Hirnantian zooplankton provinces definitions, 402

- Late Ordovician, 399–404
- Late Ordovician climate, 402–404
- Late Ordovician zooplankton provinces, 401–402
- Ordovician zooplankton, 399–401
- Sandbian palaeoceanographical boundaries, 402
- Sandbian zooplankton provinces definitions, 401–402