

Introduction to volcanism in Antarctica: 200 million years of subduction, rifting and continental break-up

John L. Smellie*¹, Kurt S. Panter² and Adelina Geyer³

¹Department of Geography, Geology and the Environment, University of Leicester, Leicester LE1 7RH, UK

²School of Earth, Environment and Society, Bowling Green State University, Bowling Green, OH 43403, USA

³Geosciences Barcelona, GEO3BCN – CSIC, Lluís Solé i Sabaris s/n, 08028 Barcelona, Spain

 JLS, 0000-0001-5537-1763; KSP, 0000-0002-0990-5880; AG, 0000-0002-8803-6504

*Correspondence: jls55@leicester.ac.uk

Antarctica has undergone several important phases of volcanism throughout its long history. It was formerly at the heart of Gondwana but, from early Jurassic time (c. 200 Ma), it commenced the prolonged process of disintegration, which resulted in the dispersal and final disposition of the southern hemisphere continents that we are familiar with today (Veevers 2012; Storey and Granot 2021). As a consequence, volcanism has been particularly important in its construction and it is geographically widespread, although mainly located within West Antarctica (Fig. 1). Its effects have frequently been felt far outside of the continent. For example, it has been a driver of global mass extinctions (Burgess *et al.* 2015; Ernst and Youbi 2017) and it has potentially driven Antarctica climatically, and by implication the world, both into and out of glacials (Bay *et al.* 2006; McConnell *et al.* 2017). Conversely, Antarctica's volcanoes may have played a pivotal role in helping Life not only to survive multiple glacial episodes during the past few tens of millions of years but to undergo species diversification on the continent in spite of the dramatic climate variations (Fraser *et al.* 2014). Eruptions from Mount Erebus also represent a significant point source of gases and aerosols to the Austral polar troposphere, including affecting the ozone layer (Boichu *et al.* 2011; Zuev *et al.* 2015). Some of Antarctica's active volcanoes also have the potential to have a significant impact on southern hemisphere aviation (Geyer *et al.* 2017). Finally, Antarctica contains the world's largest and longest-lived glaciovolcanic province. The glaciovolcanic sequences contain a detailed record of the terrestrial Antarctic ice sheet going back to nearly 30 Ma and that record is now beginning to be tapped (e.g. Smellie *et al.* 2009; Smellie and Edwards 2016; Wilch *et al.* 2021). Despite these attributes, however, volcanism in Antarctica remains terra incognita to many Earth scientists, probably in large part because of its remoteness and inaccessibility.

The only previous volume to be devoted solely to Antarctic volcanism was published 30 years ago (i.e. LeMasurier and Thomson 1990). The scope of that volume was also considerably more restricted than in this Memoir: that is, to volcanoes <c. 30 myr old. That necessitated a main focus on the generally well-preserved volcanoes erupted into the West Antarctic Rift System (WARS), one of the world's major continental rift zones. The young remnants of formerly much more widespread arc-related volcanism, and intraplate volcanism in the sub-Antarctic region, were also included. Nevertheless, nothing else existed at the time and it was, and remains, a landmark publication that inspired a generation of Earth scientists to work in Antarctica, particularly volcanologists and petrologists. As a result, numerous new investigations rapidly followed and our knowledge of Antarctica's volcanism increased profoundly, with the generation of abundant new

observations and many large datasets. However, as is always the case, much of the new knowledge has remained unpublished. A major intention of this volume is to capture that information and review it in a modern context after three decades of scientific advancement and enlightenment.

The Proterozoic and Paleozoic record of Antarctica's volcanism is patchy and comparatively poorly known (Riley *et al.* 2012; Goodge 2019). However, from the Early Jurassic (c. 200 Ma) onward, the record is much more complete and the multiple compositionally and tectonically diverse episodes of Antarctic volcanism have been intensively investigated. The volcanism can be divided into five categories (Smellie 2020): (1) Gondwana break-up volcanism (flood lavas and sills); (2) subduction-related continental margin arc, back-arc and marginal basin volcanism; (3) post-subduction slab-window basalts; (4) continental rift volcanism; and (5) intraplate volcanism of enigmatic origin. In this Memoir, the focus is on the better-preserved and much better-known record from c. 200 Ma, and each category is reviewed and assessed on a geographical basis in terms of its volcanology and eruptive palaeoenvironments; and petrology. Reviews of Antarctica's widely dispersed active volcanism, including tephrochronology (both onshore and offshore) and active subglacial volcanism, are also included. Overall, the objective of this Memoir is to review and assess the present state of knowledge of volcanism younger than c. 200 Ma right across Antarctica in all its aspects, and in the context of the interplay between the volcanism and the prevailing tectonic setting. Together with numerous geological maps, comprehensive tables of geochemical data and isotopic ages, and geophysical information, the intention is that this volume shall be the go-to resource for information on Antarctica's volcanism, the reference text for the coming decades. Our hope is that it shall act as a springboard for new proposals which, like the LeMasurier and Thomson volume, will revivify volcanic research in the region.

Volcanism in Antarctica: a brief overview

Gondwana break-up volcanism commencing at c. 190 Ma may have been driven by the effects of a large mantle plume (Storey 1995) and it is represented in Antarctica by two major voluminous volcanic provinces (Fig. 1a). One is a mafic large igneous province (LIP) that crops out throughout the Transantarctic Mountains and in Dronning Maud Land, with correlatives in South Africa, Tasmania, Australia and New Zealand (Elliot *et al.* 2021; Elliot and Fleming 2021; Luttinen 2021). The other consists of a series of felsic flare-ups that affected the entire Antarctic Peninsula and extended into southern South America (Chon Aike province: Riley and Leat 2021a, b). The mafic volcanism in Antarctica is




Fig. 1. Schematic time slices illustrating the different stages of volcanism that have affected Antarctica during the past 200 myr (modified after Smellie 2021a). The legends in diagrams (a) and (b) apply to diagrams (a)–(f), inclusive. In the Holocene depiction (g), only large volcanoes ($\geq c. 10$ km basal diameter) are shown. The diagram also includes Gausberg, a small, extinct, ultrapotassic volcanic centre 56 ka in age, to show its remarkable isolation from all other volcanic outcrops in Antarctica. NZ, New Zealand; CP, Campbell Plateau.

known as the Ferrar Supergroup but representatives in Dronning Maud Land show greater compositional affinities to coeval mafic lavas and sills in the Karoo of South Africa. The mafic LIP comprises voluminous flood lavas and thick dolerite sills. The event probably had a total volume of more than $0.5 \times 10^6 \text{ km}^3$ but it may have been emplaced in a very short time, perhaps as little as 400 kyr (Marsh 2007; Burgess *et al.* 2015). Magma in the sills is estimated to have travelled *c.* 4000 km laterally, making it the longest interpreted magma flow on Earth (Elliot *et al.* 1999; Leat 2008). The earliest eruptive phase was characterized by phreatomagmatic pyroclastic eruptions from nested maar–diatreme vent complexes collectively called phreatocauldrons (White and McClintock 2001).

By contrast, the felsic Chon Aike province was erupted in three major pulses, individually *c.* 6–10 myr in duration, between 189 and 153 Ma (Pankhurst *et al.* 2000; Riley *et al.* 2001; Riley and Leat 2021a, b). Each corresponds to a predominantly explosive volcanic flare-up (*sensu* Paterson and Ducea 2015) resulting mainly in the eruption of rhyolite ignimbrites. A prominent Pacific-ward progression observed in the ages of the volcanism has been linked to the impact and sublithospheric melting effects of the large spreading head of the same plume responsible for the Ferrar–Karoo LIP. An arc-like (subduction-influenced) mantle source, rather than a plume, has also been postulated for the mafic LIP (Choi *et al.* 2019; Elliot and Fleming 2021; Panter 2021).

Coincident with the break-up magmatism, the Pacific margin of Gondwana was already the locus of a major long-lived continental magmatic arc, the products of which are today widely preserved throughout the Antarctic Peninsula (Fig. 1b–e) (Leat and Riley 2021a, b). The volcanism in the arc, and its plutonic equivalents, have been extensively dated, revealing that the activity probably took place as a series of magmatic flare-ups and lulls (Riley *et al.* 2018). This is particularly evident in the South Shetland Islands, which is the most intensively dated arc-related region (Haase *et al.* 2012; Leat and Riley 2021a). The locus of active arc volcanism in Palmer Land and much of Graham Land moved into the forearc region during the Late Cretaceous, probably due to orthogonal subduction coupled with enhanced time-integrated growth of an extensive accretionary complex. By contrast, the volcanism migrated obliquely away from the trench in the South Shetland Islands possibly due to a highly oblique subduction vector causing much more limited accretion and possibly forearc erosion (Smellie 2021a). The magmatic activity shut down progressively in a clockwise direction, commencing in Marie Byrd Land in the mid-Cretaceous (Larter *et al.* 2002). Subduction is active today (at a very slow rate) only at the northern tip of the Antarctic Peninsula, where a small ensialic marginal basin populated with numerous submarine volcanic centres opened up in response to plate boundary forces, including slab rollback (Haase and Beier 2021; Smellie 2021b). The marginal basin includes Deception Island, one of Antarctica's most active volcanoes (Geyer *et al.* 2021), which underwent a major caldera collapse eruption *c.* 4 kyr ago that dispersed ash more than 4000 km in an arcuate swathe across the Scotia Sea and much of East Antarctica (Antoniades *et al.* 2018). Coincidentally, from *c.* 12 Ma an extensive back-arc mafic alkaline volcanic field developed to the rear of Graham Land in the James Ross Island region (Fig. 1f) (Haase and Beier 2021; Smellie 2021b). It is dominated by the very large shield volcano of Mount Haddington, which is predominantly glaciovolcanic and contains an unrivalled record of the Antarctic Peninsula Ice Sheet (Smellie *et al.* 2008).

Following the progressive shut down of subduction, 'windows' opened up in the downgoing oceanic slab, allowing ingress and decompression melting of mantle unaffected by

subduction metasomatism. As a consequence, several small monogenetic alkaline volcanic fields were constructed from *c.* 7.5 Ma, mainly along the flanks of the Antarctic Peninsula (Fig. 1f) (Hole *et al.* 1995). The outcrops are overwhelmingly glaciovolcanic and, like Mount Haddington, they also preserve a uniquely valuable record of the Antarctic Peninsula Ice Sheet, as well as influencing our understanding of glaciovolcanism (Smellie and Edwards 2016; Smellie and Hole 2021). New research suggests that melting of the downgoing slab may be involved in the generation of the slab-window basalts (Hole 2021).

Over a period of a few million years in the Cretaceous, Antarctica's magmatism shifted from arc-related to rift-related during the latest stage of Gondwana break-up that separated Zealandia from Antarctica (*c.* 90–83 Ma). Rare Late Cretaceous alkaline intraplate magmatism is found in southern Zealandia (Weaver and Smith 1989), including HIMU-like ocean island basalt (OIB) (Panter *et al.* 2006; Hoernle *et al.* 2020), and mafic dykes are present along the Ruppert and Hobbs coasts of Marie Byrd Land (Storey *et al.* 1999). However, the episode was largely amagmatic. The extension also probably caused widespread topographical lowering until much of the region in Marie Byrd Land (and coeval terrain in New Zealand) subsided down to, or close to, sea level (LeMasurier and Landis 1996).

Renewed extension during the Cenozoic resulted in the creation of the WARS, a very large continental rift characterized by widespread alkaline volcanism (Fig. 1d–f) (Siddoway 2008; Smellie and Martin 2021; Smellie and Rocchi 2021; Wilch *et al.* 2021). The rift contains numerous large and small volcanoes with basalt–trachyte, phonolite and rhyolite compositions (Martin *et al.* 2021; Panter *et al.* 2021b; Rocchi and Smellie 2021). Remote sensing studies have also suggested that numerous additional volcanic centres may be widespread beneath the West Antarctic Ice Sheet, including several that may be active (Behrendt *et al.* 1994; van Wyk de Vries *et al.* 2018; Quartini *et al.* 2021). The origins of the volcanism are disputed, and variable roles have been inferred for deep mantle plumes and shallow thermal anomalies with associated edge flow (LeMasurier and Landis 1996; Rocchi *et al.* 2005; Martin *et al.* 2021; Panter *et al.* 2021b; Rocchi and Smellie 2021). They are all linked within the broad concept of a diffuse alkaline magmatic province (DAMP), which also includes Late Cretaceous and Cenozoic volcanism in eastern Australia, Tasmania and Zealandia (Finn *et al.* 2005). The striking compositional commonality throughout the DAMP region has been explained by melting of a mantle source component inferred to underlie the entire region, with the characteristics of HIMU-like OIB. The mantle reservoir may have been emplaced as a large plume head, of late Cretaceous age or older, or lithosphere that has been metasomatized, or both sources exist (Panter 2021). Like alkaline volcanism in the Antarctic Peninsula, the volcanism in the WARS is also predominantly glaciovolcanic, and it has also provided uniquely important information on the development of the West and East Antarctic ice sheets (Smellie and Rocchi 2021; Wilch *et al.* 2021).

The tectonic setting of some of Antarctica's volcanoes is enigmatic, however. They include at least three small monogenetic phreatomagmatic edifices in the southern Transantarctic Mountains (upper Scott Glacier) and beneath the East Antarctic Ice Sheet, less than 300 km from the South Pole (Fig. 1e) (Smellie *et al.* 2021), and Gaussberg, which is an isolated pillow volcano on the East Antarctic coast far from any other expression of volcanism on the continent (Fig. 1g) (Smellie and Collerson 2021). The upper Scott Glacier and nearby subglacial centres are outside of the WARS and they are believed to have formed in response to the detachment and sinking of lithosphere into the convecting mantle beneath

the East Antarctic Craton (Panter *et al.* 2021a). The composition of Gausberg is unique in Antarctica and very rare worldwide, being formed of ulrapotassic lamproite. Its origin is probably linked to a small deep-sourced mantle plume, distinct from the large Kerguelen plume, that incorporated a component derived from ancient subducted sediment (Murphy *et al.* 2002).

Antarctica also contains several large active volcanoes (Geyer 2021) (Fig. 1g). Only two have been observed in eruption (Mount Erebus and Deception Island); both have been intensively investigated (Oppenheimer and Kyle 2008; Geyer *et al.* 2021; Sims *et al.* 2021). Mount Erebus also hosts the world's only semi-permanent phonolite lava lake. The presence of relict heat (Mount Berlin, Mount Melbourne and Mount Rittmann) and abundant englacial and marine tephra sourced in Mount Takahe, Mount Berlin, Mount Waesche, Mount Rittmann and, possibly, The Pleiades indicate that many others were active in recent geological time (<10 ka: Lee *et al.* 2019; Dunbar *et al.* 2021; Gambino *et al.* 2021; Narcisi and Petit 2021; Di Roberto *et al.* 2021). Three of the volcanoes are, or have been, monitored (Deception Island, Mount Erebus and Mount Melbourne) but only one has published hazard and risk assessments (Deception Island: Bartolini *et al.* 2014; Pedrazzi *et al.* 2018; Geyer *et al.* 2021).

Acknowledgements The authors gratefully acknowledge the collaborations and conversations with colleagues, too many to mention, over the many years that we have been actively engaged in Antarctic research. We are also grateful for the support provided by the various national funding agencies, without which none of our studies could have been implemented.

Author contributions JLS: conceptualization (lead), writing – original draft (lead); KSP: conceptualization (supporting); AG: conceptualization (supporting).

Funding This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Data availability Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

References

- Antoniades, D., Giralt, S. *et al.* 2018. The timing and widespread effects of the largest Holocene volcanic eruption in Antarctica. *Scientific Reports*, **8**, 17279, <https://doi.org/10.1038/s41598-018-35460-x>
- Bartolini, S., Geyer, A., Martí, J., Pedrazzi, D. and Aguirre-Díaz, G. 2014. Volcanic hazard on Deception Island (South Shetland Islands, Antarctica). *Journal of Volcanology and Geothermal Research*, **285**, 150–168, <https://doi.org/10.1016/j.jvolgeores.2014.08.009>
- Bay, R.C., Bramall, N.E., Price, P.B., Clow, G.D., Hawley, R.L., Udisti, R. and Castellano, E. 2006. Globally synchronous ice core volcanic tracers and abrupt cooling during the last glacial period. *Journal of Geophysical Research: Atmospheres*, **111**, D11108, <https://doi.org/10.1029/2005JD006306>
- Behrendt, J.C., Blankenship, D.D., Finn, C.A., Bell, R.E., Sweeney, R.E., Hodge, S.R. and Brozena, J.M. 1994. CASERTZ aeromagnetic data reveal late Cenozoic flood basalts(?) in the West Antarctic rift system. *Geology*, **22**, 527–530, [https://doi.org/10.1130/0091-7613\(1994\)022<0527:CADRLC>2.3.CO;2](https://doi.org/10.1130/0091-7613(1994)022<0527:CADRLC>2.3.CO;2)
- Boichu, M., Oppenheimer, C., Roberts, T.J., Tsanev, V. and Kyle, P. 2011. On bromine, nitrogen oxides and ozone depletion in the tropospheric plume of Erebus volcano (Antarctica). *Atmospheric Environment*, **45**, 3856–3866, <https://doi.org/10.1016/j.atmosenv.2011.03.027>
- Burgess, S.D., Bowring, S.A., Fleming, T.H. and Elliot, D.H. 2015. High precision geochronology links the Ferrar Large Igneous Province with early Jurassic ocean anoxia and biotic crisis. *Earth and Planetary Science Letters*, **415**, 90–99, <https://doi.org/10.1016/j.epsl.2015.01.037>
- Choi, S.H., Mukasa, S.B., Ravizza, G., Fleming, T.H., Marsh, B.D. and Bédard, J.H.J. 2019. Fossil subduction zone origin for magmas in the Ferrar Large Igneous Province, Antarctica: evidence from PGE and OS isotope systematics in the Basement Sill of the McMurdo Dry Valleys. *Earth and Planetary Science Letters*, **506**, 507–519, <https://doi.org/10.1016/j.epsl.2018.11.027>
- Di Roberto, A., del Carlo, P. and Pompilio, M. 2021. Marine record of Antarctic volcanism from drill cores. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-49>
- Dunbar, N.W., Iverson, N.A., Smellie, J.L., McIntosh, W.C., Zimmerer, M.J. and Kyle, P.R. 2021. Active volcanoes in Marie Byrd Land. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-29>
- Elliot, D.H. and Fleming, T.H. 2021. Ferrar Large Igneous Province: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-39>
- Elliot, D.H., Fleming, T.W., Kyle, P.R. and Foland, K.A. 1999. Long-distance transport of magmatism the Jurassic Ferrar Large Igneous Province, Antarctica. *Earth and Planetary Science Letters*, **167**, 89–104, [https://doi.org/10.1016/S0012-821X\(99\)00023-0](https://doi.org/10.1016/S0012-821X(99)00023-0)
- Elliot, D.H., White, J.D.L. and Fleming, T.H. 2021. Ferrar Large Igneous Province: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-44>
- Ernst, R.E. and Youbi, N. 2017. How large igneous provinces affect global climate, sometimes cause mass extinctions, and represent natural markers in the geological record. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **478**, 30–52, <https://doi.org/10.1016/j.palaeo.2017.03.014>
- Finn, C.A., Müller, R.D. and Panter, K.S. 2005. A Cenozoic diffuse alkaline magmatic province (DAMP) in the southwest Pacific without rift or plume origin. *Geochemistry, Geophysics, Geosystems*, **6**, Q02005, <https://doi.org/10.1029/2004GC000723>
- Fraser, C.I., Terauds, A., Smellie, J., Convey, P. and Chown, S. 2014. Geothermal activity helps life survive glacial cycles. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 5634–5639, <https://doi.org/10.1073/pnas.1321437111>
- Gambino, S., Armienti, P. *et al.* 2021. Mount Melbourne and Mount Rittmann. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-43>
- Geyer, A. 2021. Antarctic volcanism: active volcanism overview. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2020-12>
- Geyer, A., Martí, A., Giralt, S. and Folch, A. 2017. Potential ash impact from Antarctic volcanoes: Insights from Deception Island's most recent eruption. *Scientific Reports*, **7**, 16534, <https://doi.org/10.1038/s41598-017-16630-9>
- Geyer, A., Pedrazzi, D. *et al.* 2021. Deception Island. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-56>
- Goodge, J.W. 2019. Geological and tectonic evolution of the Transantarctic Mountains, from ancient craton to recent enigma. *Gondwana Research*, **80**, 50–122, <https://doi.org/10.1016/j.gr.2019.11.001>
- Haase, K.M. and Beier, C. 2021. Bransfield Strait and James Ross Island: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-37>
- Haase, K.M., Beier, C., Fretzdorff, S., Smellie, J.L. and Garbe-Schönberg, D. 2012. Magmatic evolution of the South Shetland Islands, Antarctica, and implications for continental crust formation. *Contributions to Mineralogy and Petrology*, **163**, 1103–1119, <https://doi.org/10.1007/s00410-012-0719-7>
- Hoernle, K., Timm, C. *et al.* 2020. Late Cretaceous (99–69 Ma) basaltic intraplate volcanism on and around Zealandia: Tracing

- upper mantle geodynamics from Hikurangi Plateau collision to Gondwana breakup and beyond. *Earth and Planetary Science Letters*, **529**, 115864, <https://doi.org/10.1016/j.epsl.2019.115864>
- Hole, M.J. 2021. Antarctic Peninsula: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-40>
- Hole, M.J., Saunders, A.D., Rogers, G. and Sykes, M.A. 1995. The relationship between alkaline magmatism, lithospheric extension and slab window formation along continental destructive plate margins. *Geological Society, London, Special Publications*, **81**, 265–285, <https://doi.org/10.1144/GSL.SP.1994.081.01.15>
- Larter, R.D., Cunningham, A.P., Barker, P.F., Gohl, K. and Nitsche, F.O. 2002. Tectonic evolution of the Pacific margin of Antarctica 1. Late Cretaceous tectonic reconstructions. *Journal of Geophysical Research: Solid Earth*, **107**, 2345, <https://doi.org/10.1029/2000JB000052>
- Leat, P.T. 2008. On the long-distance transport of Ferrar magmas. *Geological Society, London, Special Publications*, **302**, 45–61, <https://doi.org/10.1144/SP302.4>
- Leat, P.T. and Riley, T.R. 2021a. Antarctic Peninsula and South Shetland Islands: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-52>
- Leat, P.T. and Riley, T.R. 2021b. Antarctic Peninsula and South Shetland Islands: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-68>
- Lee, M.J., Kyle, P.R., Iverson, N.A., Lee, J.I. and Han, Y. 2019. Rittmann volcano, Antarctica as the source of a widespread 1252 ± 2 CE tephra layer in Antarctica ice. *Earth and Planetary Science Letters*, **521**, 169–176, <https://doi.org/10.1016/j.epsl.2019.06.002>
- LeMasurier, W.E. and Landis, C.A. 1996. Mantle-plume activity recorded by low-relief erosion surfaces in West Antarctica and New Zealand. *Geological Society of America Bulletin*, **108**, 1450–1466, [https://doi.org/10.1130/0016-7606\(1996\)108<1450:MPARBL>2.3.CO;2](https://doi.org/10.1130/0016-7606(1996)108<1450:MPARBL>2.3.CO;2)
- LeMasurier, W.E. and Thomson, J.W. (eds) 1990. *Volcanism of the Antarctic Plate and Southern Oceans*. American Geophysical Union Antarctic Research Series, **48**.
- Luttinen, A.V. 2021. Dronning Maud Land Jurassic volcanism: volcanology and petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-89>
- Marsh, B.D. 2007. Magmatism, magma, and magma chambers. *Treatise on Geophysics*, **6**, 276–333.
- Martin, A.P., Cooper, A.F., Price, R.C., Kyle, P.R. and Gamble, J.A. 2021. Erebus Volcanic Province: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-80>
- McConnell, J.R., Burke, A. *et al.* 2017. Synchronous volcanic eruptions and abrupt climate change similar to 17.7 ka plausibly linked by stratospheric ozone depletion. *Proceedings of the National Academy of Sciences of the United States of America*, **114**, 10 035–10 040, <https://doi.org/10.1073/pnas.1705595114>
- Murphy, D.T., Collerson, K.D. and Kamber, B.S. 2002. Lamproites from Gaussberg, Antarctica: possible transition zone melts of Archaean subducted sediments. *Journal of Petrology*, **43**, 981–1001, <https://doi.org/10.1093/ptrology/43.6.981>
- Narcisi, B. and Petit, J.R. 2021. Englacial tephra of East Antarctica. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-86>
- Oppenheimer, C. and Kyle, P.R. (eds) 2008. Volcanology of Erebus volcano, Antarctica. *Journal of Volcanology and Geothermal Research*, **177**, 531–754, <https://doi.org/10.1016/j.jvolgeores.2008.08.016>
- Pankhurst, R.J., Riley, T.R., Fanning, C.M. and Kelley, S.P. 2000. Episodic silicic volcanism in Patagonia and the Antarctic Peninsula: Chronology of magmatism associated with the break-up of Gondwana. *Journal of Petrology*, **41**, 605–625, <https://doi.org/10.1093/ptrology/41.5.605>
- Panter, K.S. 2021. Antarctic volcanism: petrology and tectonomagmatic overview. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2020-10>
- Panter, K.S., Blusztajn, J., Hart, S.R., Kyle, P.R., Esser, R. and McIntosh, W.C. 2006. The origin of HIMU in the SW Pacific: Evidence from intraplate volcanism in southern New Zealand and subantarctic islands. *Journal of Petrology*, **47**, 1673–1704, <https://doi.org/10.1093/ptrology/egl024>
- Panter, K.S., Reindel, J. and Smellie, J.L. 2021a. Mount Early and Sheridan Bluff: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-2>
- Panter, K.S., Wilch, T.I., Smellie, J.L., Kyle, P.R. and McIntosh, W.C. 2021b. Marie Byrd Land and Ellsworth Land: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-50>
- Paterson, S.R. and Ducea, M.N. 2015. Arc magmatic tempos: gathering the evidence. *Elements*, **11**, 91–98, <https://doi.org/10.2113/gselements.11.2.91>
- Pedraza, D., Németh, K., Geyer, A., Álvarez-Valero, A.M., Aguirre-Díaz, G. and Bartolini, S. 2018. Historic hydrovolcanism at Deception Island (Antarctica): implications for eruption hazards. *Bulletin of Volcanology*, **80**, 11, <https://doi.org/10.1007/s00445-017-1186-9>
- Quartini, E., Blankenship, D.D. and Young, D.A. 2021. Active subglacial volcanism in West Antarctica. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-3>
- Riley, T.R. and Leat, P.T. 2021a. Palmer Land and Graham Land volcanic groups (Antarctic Peninsula): volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-36>
- Riley, T.R. and Leat, P.T. 2021b. Palmer Land and Graham Land volcanic groups (Antarctic Peninsula): petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-51>
- Riley, T.R., Leat, P.T., Pankhurst, R.J. and Harris, C. 2001. Origins of large volume rhyolitic volcanism in the Antarctic Peninsula and Patagonia by crustal melting. *Journal of Petrology*, **42**, 1043–1065, <https://doi.org/10.1093/ptrology/42.6.1043>
- Riley, T.R., Flowerdew, M.J. and Whitehouse, M.J. 2012. U–Pb ion-microprobe zircon geochronology from the basement inliers of eastern Graham Land, Antarctic Peninsula. *Journal of the Geological Society, London*, **169**, 381–393, <https://doi.org/10.1144/0016-76492011-142>
- Riley, T.R., Burton-Johnson, A., Flowerdew, M.J. and Whitehouse, M.J. 2018. Episodicity within a mid-Cretaceous magmatic flare-up in West Antarctica: U–Pb ages of the Lassiter Coast intrusive suite, Antarctic Peninsula, and correlations along the Gondwana margin. *Geological Society of America Bulletin*, **130**, 1177–1196, <https://doi.org/10.1130/B31800.1>
- Rocchi, S. and Smellie, J.L. 2021. Northern Victoria Land: petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-19>
- Rocchi, S., Armienti, P. and Di Vincenzo, G. 2005. No plume, no rift magmatism in the West Antarctic Rift. *United States Geological Survey Professional Papers*, **388**, 435–447.
- Siddoway, C. 2008. Tectonics of the West Antarctic rift system: New light on the history and dynamics of distributed intracontinental extension (invited paper). In: Cooper, A.K., Barrett, P.J. *et al.* (eds) *Antarctica: A Keystone in a Changing World. Proceedings of the 10th International Symposium on Antarctic Earth Sciences*. The National Academies Press, Washington, DC, 91–114.
- Sims, K.W.W., Aster, R. *et al.* 2021. Mount Erebus. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-8>
- Smellie, J.L. 2020. The role of volcanism in the making of Antarctica. In: Oliva, M. and Ruiz-Fernández, J. (eds) *Past Antarctica. Paleoclimatology and Climate Change*. Academic Press, London, 69–88.
- Smellie, J.L. 2021a. Antarctic volcanism: volcanology and palaeoenvironmental overview. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2020-1>
- Smellie, J.L. 2021b. Bransfield Strait and James Ross Island: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-58>
- Smellie, J.L. and Collerson, K.D. 2021. Gaussberg: volcanology and petrology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-85>
- Smellie, J.L. and Edwards, B.E. 2016. *Glaciovolcanism on Earth and Mars: Products, Processes and Palaeoenvironmental Significance*. Cambridge University Press, Cambridge, UK.

- Smellie, J.L. and Hole, M.J. 2021. Antarctic Peninsula: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-59>
- Smellie, J.L. and Martin, A.P. 2021. Erebus Volcanic Province: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-62>
- Smellie, J.L. and Rocchi, S. 2021. Northern Victoria Land: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-60>
- Smellie, J.L., Johnson, J.S., McIntosh, W.C., Esser, R., Gudmundsson, M.T., Hambrey, M.J. and van Wyk de Vries, B. 2008. Six million years of glacial history recorded in the James Ross Island Volcanic Group, Antarctic Peninsula. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **260**, 122–148, <https://doi.org/10.1016/j.palaeo.2007.08.011>
- Smellie, J.L., Haywood, A.M., Hillenbrand, C.-D., Lunt, D.J. and Valdes, P.J. 2009. Nature of the Antarctic Peninsula Ice Sheet during the Pliocene: Geological evidence and modelling results compared. *Earth-Science Reviews*, **94**, 79–94, <https://doi.org/10.1016/j.earscirev.2009.03.005>
- Smellie, J.L., Panter, K.S. and Reindel, J. 2021. Mount Early and Sheridan Bluff: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-61>
- Storey, B.C. 1995. The role of mantle plumes in continental breakup: case histories from Gondwanaland. *Nature*, **377**, 301–308, <https://doi.org/10.1038/377301a0>
- Storey, B.C. and Granot, R. 2021. Tectonic history of Antarctica over the past 200 million years. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2018-38>
- Storey, B.C., Leat, P.T., Weaver, S.D., Pankhurst, R.J., Bradshaw, J.D. and Kelley, S. 1999. Mantle plumes and Antarctica–New Zealand rifting: evidence from mid-Cretaceous mafic dykes. *Journal of the Geological Society, London*, **156**, 659–671, <https://doi.org/10.1144/gsjgs.156.4.0659>
- van Wyk de Vries, M., Bingham, R.G. and Hein, A.S. 2018. A new volcanic province: an inventory of subglacial volcanoes in West Antarctica. *Geological Society, London, Special Publications*, **461**, 231–248, <https://doi.org/10.1144/SP461.7>
- Veevers, J.J. 2012. Reconstructions before rifting and drifting reveal the geological connections between Antarctica and its conjugates in Gondwanaland. *Earth-Science Reviews*, **111**, 249–318, <https://doi.org/10.1016/j.earscirev.2011.11.009>
- Weaver, S.D. and Smith, I.E.M. 1989. New Zealand intraplate volcanism. In: Johnson, R.W., Knutson, J. and Taylor, S.R. (eds) *Intraplate Volcanism in Eastern Australia and New Zealand*. Cambridge University Press, Cambridge, UK, 157–188.
- White, J.D.L. and McClintock, M.K. 2001. Immense vent complex marks flood-basalt eruption in a wet, failed rift: Coombs Hills, Antarctica. *Geology*, **29**, 935–938, [https://doi.org/10.1130/0091-7613\(2001\)029<0935:IVCMFB>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0935:IVCMFB>2.0.CO;2)
- Wilch, T.I., McIntosh, W.C. and Panter, K.S. 2021. Marie Byrd Land and Ellsworth Land: volcanology. *Geological Society, London, Memoirs*, **55**, <https://doi.org/10.1144/M55-2019-39>
- Zuev, V.V., Zueva, N.E., Savelieva, E.S. and Gerasimov, V.V. 2015. The Antarctic ozone depletion caused by Erebus volcano gas emissions. *Atmospheric Environment*, **122**, 393–399, <https://doi.org/10.1016/j.atmosenv.2015.10.005>